arXiv:1301.6451v2 [math.CO] 4 Feb 2013

On symmetry groups of oriented matroids

Hiroyuki Miyata
Graduate School of Information Science, Tohoku University
Aramaki Aza Aoba, Aoba-ku, Sendai, 980-8579 Japan
`hmiyata@dais.is.tohoku.ac.jp`

February 5, 2013

Abstract

Symmetry of hyperplane arrangements, point configurations and polytopes have been of great interests among many people, and have been studied extensively for a long time. On the other hand, symmetry of oriented matroids, combinatorial abstractions of them, is not understood so well.

In this paper, we aim to obtain a better understanding of symmetry of oriented matroids. First, we focus on matroids and give a general construction that generates a 3-dimensional point configuration with a matroidal symmetry that cannot be realized as a geometric symmetry. The construction is motivated by the fact that every non-trivial rotation in the 2-dimensional Euclidean space has a unique fixed point but there is no corresponding property for matroids. The construction suggests that the lack of the fixed point theorem generates a big gap between matroidal symmetries and geometric symmetries of point configurations. Motivated by this insight, we investigate some fixed point properties for oriented matroids. We develop some fixed point theorems for oriented matroids and then make classification of rotational symmetry groups of simple acyclic oriented matroids of rank 4 as an application.

1 Introduction

Symmetry of geometric structures such as polytopes, hyperplane arrangements and point configurations has been paid great interests for a long time and there are rich theories for them (see [8, 14], for example). A natural question is whether symmetry of *oriented matroids* and *matroids*, ones of the most well-studied combinatorial abstractions of polytopes, hyperplane arrangements and point configurations, have similar properties or not. However, to the best of the author's knowledge, it is not understood so well. In this paper, we aim to contribute to a better understanding of symmetry of matroids and oriented matroids.

First, we investigate a gap between *geometric symmetry* and *combinatorial symmetry* of point configurations. Geometric symmetry is symmetry defined by affine invariance. By combinatorial symmetry, we mean symmetry of the underlying combinatorial structures such as the face lattices of polytopes, and the matroids or the oriented matroids of point configurations. Every geometric symmetry induces a combinatorial symmetry, but the converse is not always true. There are considerable studies on this topic since it is an important step to understand when combinatorial symmetry cannot be realized geometrically, in order to analyze symmetry of matroids and oriented matroids. Here, we review studies on gaps between combinatorial symmetry and geometric symmetry of related structures.

In 1971, Mani [10] proved that every combinatorial symmetry (face-lattice symmetry) can be realized geometrically for 3-polytopes. Then Perles showed that the same holds for d-polytopes with $d+3$ vertices [7, p.120]. After that, it had been a big question whether every combinatorial symmetry (face-lattice symmetry) can be realized geometrically for every polytope. Bokowski, Ewald and Kleinschmidt [4] resolved the question by proving that there is a 4-polytope with 10 vertices with a non-realizable combinatorial symmetry (face-lattice symmetry). For point configurations, Shor [13] constructed an example of a 2-dimensional configuration of 64 points with a non-realizable combinatorial symmetry (oriented-matroid symmetry) and Richter-Gebert [12] proposed an example of a 2-dimensional configuration of 14 points with a non-realizable combinatorial symmetry (oriented-matroid symmetry). In 2006, Paffenholz [11] constructed a two-parameter infinite family of 4-polytopes with non-realizable combinatorial symmetries (face-lattice symmetries).

Our contribution:

First, we study a gap between combinatorial symmetry (matroidal symmetry) and geometric symmetry of point configurations. We consider a general method to construct 3-dimensioinal point configurations with non-realizable matroidal symmetries. Similarly to the constructions of polytopes with non-realizable combinatorial symmetries by Bokowski, Ewald and Kleinschmidt [4] and Paffenholz [11], the key tool for our construction is the fixed point theorem, which asserts that every non-trivial rotation in the 2-dimensional Euclidean space has a unique fixed point. There is no corresponding property for matroids and it makes the gaps between geometric symmetry and matroidal symmetry. Motivated by this insight, we investigate some fixed point properties for oriented matroids. We study rotational symmetry groups of simple acyclic oriented matroids of rank 4 using obtained theorems. In particular, it is proved that all possible rotational symmetry groups of acyclic simple oriented matroids of rank 4 are covered by cyclic groups C_n $(n \geq 1)$, dihedral groups D_{2n} $(n \geq 1)$, the alternating group A_4, the symmetry group S_4 and the alternating group A_5. The result completely coincides with the classification of rotational symmetry groups of 3-dimensional point configurations.

Organization of the paper:

In Section 2, we explain some terminologies on oriented matroids. Section 3 is dedicated to studying symmetry of matroids. We give a general construction of 3-dimensional point configuratoins with non-realizable matroidal symmetries. Motivated by insights in Section 3, we study fixed point theorems for oriented matroids in Section 4. In Section 5, symmetry groups of simple oriented matroids of rank 3 are investigated. Based on results in Sections 4 and 5, we classify rotational symmetry groups of acyclic simple oriented matroids of rank 4 in Section 6. In Section 7, we make a conclusion of the paper.

2 Preliminaries

In this paper, we assume that the reader is familiar with oriented matroids. Here, we make a brief introduction. For details on oriented matroids, see [2].

2.1 Definitions on oriented matroids

Oriented matroids have various equivalent axiom systems. First, we explain the *chirotope axioms.*

Definition 2.1 (Chirotope axioms)
Let E be a finite set and $r \geq 1$ an integer. A *chirotope of rank* r *on* E is a mapping $\chi : E^r \to \{+1, -1, 0\}$ which satisfies the following properties for any $i_1, \dots, i_r, j_1, \dots, j_r \in E$.

$(B1)$ χ is not identically zero.

$(B2)$ $\chi(i_{\sigma(1)}, \dots, i_{\sigma(r)}) = \mathrm{sgn}(\sigma)\chi(i_1, \dots, i_r)$ for all $i_1, \dots, i_r \in E$ and any permutation σ.

$(B3)$ If $\chi(j_s, i_2, \dots, i_r) \cdot \chi(j_1, \dots, j_{s-1}, i_1, j_{s+1}, \dots, j_r) \geq 0$ for all $s = 1, \dots, r,$ then $\chi(i_1, \dots, i_r) \cdot \chi(j_1, \dots, j_r) \geq 0.$

A pair $(E, \{\chi, -\chi\})$ is called an *oriented matroid* of rank r on a ground set E. By $rank(\mathcal{M})$, we denote the rank of $\mathcal{M}$. The *underlying matroid* $\underline{\mathcal{M}}$ of $\mathcal{M}$ is defined as the matroid on the ground set E with the set of bases $\{\{i_1, \dots, i_r\} \mid \chi(i_1, \dots, i_r) \neq 0\}$. A subset $F \subset E$ is said to be a flat of $\mathcal{M}$ if F is a flat of $\underline{\mathcal{M}}$. For $A \subset E$, we denote by $Flat(A)$ the minimal flat of $\mathcal{M}$ that contains A.

For a d-dimensional point configuration $P = \{\boldsymbol{p}_1, \ldots, \boldsymbol{p}_n\}$, the *associated oriented matroid* is defined as follows. Let $\boldsymbol{v}_1, \ldots, \boldsymbol{v}_n \in \mathbb{R}^{d+1}$ is the associated vector configuration i.e.,

$$\boldsymbol{v}_1 := \begin{pmatrix} \boldsymbol{p}_1 \\ 1 \end{pmatrix}, \ldots, \boldsymbol{v}_n := \begin{pmatrix} \boldsymbol{p}_n \\ 1 \end{pmatrix}.$$

Let us define a map $\chi_P : \{1, \ldots, n\}^{d+1} \to \{0, +, -\}$ by

$$\chi_P(i_1, \ldots, i_{d+1}) := \mathrm{sign}(\det(\boldsymbol{v}_{i_1}, \ldots, \boldsymbol{v}_{i_{d+1}})) \text{ for } 1 \leq i_1, \ldots, i_{d+1} \leq n.$$

The oriented matroid associated to P is defined as the pair $(\{1, \ldots, n\}, \{\chi_P, -\chi_P\})$.

An oriented matroid can also be specified by a collection of *covectors*. In the following axiom system, we use the following notation. For sign vectors $X, Y \in \{0, +, -\}^E$, a sign vector $X \circ Y \in \{0, +, -\}^E$ is defined as follows.

$$(X \circ Y)(e) := \begin{cases} X(e) & \text{if } X(e) \neq 0, \\ Y(e) & \text{otherwise.} \end{cases}$$

We also use the following notations.

$$S(X, Y) := \{e \in E \mid X(e) = -Y(e) (\neq 0)\}.$$

$$X \geq Y \Leftrightarrow X(e) = Y(e) \text{ for all } e \in E \text{ such that } Y(e) \neq 0.$$

Definition 2.2 (Covector axioms)
An element of a set $\mathcal{V}^* \subseteq \{0, +, -\}^E$ of sign vectors satisfying the following axioms is called a *covector*.

$(V0)$ $(0 \ldots 0) \in \mathcal{V}^*$.

$(V1)$ $X \in \mathcal{V}^*$ implies $-X \in \mathcal{V}^*$.

$(V2)$ For $X, Y \in \mathcal{V}^*$, $X \circ Y \in \mathcal{V}^*$.

$(V3)$ For $X, Y \in \mathcal{V}^*$ and $e_0 \in S(X, Y)$, there exists a covector $Z \in \mathcal{V}^*$ such that $Z(e_0) = 0$ and $Z(e) = (X \circ Y)(e)$ for all $e \in E \setminus S(X, Y)$.

(vector elimination)

The axiom $(V3)$ can be replaced by the following axiom.

$(V3^f)$ For $X, Y \in \mathcal{V}^*$ and $U \subseteq S(X, Y)$, there exists a covector $Z \in \mathcal{V}^*$ and $u \in U$ such that $Z(u) = 0$, $Y|_U \geq Z|_U$ and $Z(e) = (X \circ Y)(e)$ for all $e \in E \setminus S(X, Y)$.

This operation is called *conformal elimination.*

It is sufficient to consider minimal covectors to specify an oriented matroid. An element of the set $\mathcal{C}^*$ defined as follows is called a *cocircuit.*

$$\mathcal{C}^* := \{X \in \mathcal{V}^* \mid X \not> Y \text{ for all } Y \in \mathcal{V}^* \setminus \{0\}\}.$$

The set $\mathcal{C}^*$ can be characterized by a simple axiom system, called *cocircuit axioms.* There are many other equivalent data, such as *vectors* and *circuits*, that can be used to specify oriented matroids. For details, see [2, Chapter 3].

For an oriented matroid $\mathcal{M} = (E, \mathcal{V}^*)$ specified by the set of covectors $\mathcal{V}^*$ and a subset $A \subseteq E$, the restriction $\mathcal{M}|_A$ is an oriented matroid $(A, \mathcal{V}^*|_A)$, where $\mathcal{V}^*|_A := \{V|_A \mid V \in \mathcal{V}^*\}$. We sometimes write $rank(A)$ instead of $rank(\mathcal{M}|_A)$ when there is no confusion.

An element $e \in E$ is called a *loop* if $X(e) = 0$ for all $X \in \mathcal{V}^*$. An oriented matroid is said to be *loopless* if it has no loops. If $X(e) = X(f)$ (resp. $X(e) = -X(f)$) for all $X \in \mathcal{V}^*$, e and f are said to be *parallel* (resp. *antiparallel*). An oriented matroid is said to be *simple* if it has neigther loops, distinct parallel elements nor distinct antiparallel elements. If an oriented matroid contains the positive covector $(+ + \cdots +)$, it is said to be *acyclic*. For an acyclic oriented matroid $\mathcal{M}$ on a ground set E, an element $e \in E$ is called an *extreme point* of $\mathcal{M}$ if there is the covector X such that $X(e) = 0$ and $X(f) = +$ for all $f \in E \setminus \{e\}$. An acyclic oriented matroid $\mathcal{M}$ is said to be *a matroid polytope* if all elements of $\mathcal{M}$ are extreme points.

In the rank 3 case, every matroid polytope is known to be *relabeling equivalent* to an *alternating matroid.*

Definition 2.3

- Oriented matroids $\mathcal{M} = (E, \mathcal{V}^*)$ and $\mathcal{M}' = (E', \mathcal{V}'^*)$ are said to be *relabeling equivalent* if there exists a bijection $\phi : E \to E'$ such that $X \in \mathcal{V}^* \Leftrightarrow \phi(X) \in \mathcal{V}'^*$.
- On the other hand, $\mathcal{M}$ and $\mathcal{M}'$ are said to be *reorientation equivalent* if ${}_{-A}\mathcal{M}$ and $\mathcal{M}'$ are relabeling equivalent for some $A \subseteq E$. Here, ${}_{-A}\mathcal{M}$ is the oriented matroid specified by the collection of covectors $\{{}_{-A}X \mid X \in \mathcal{V}^*\}$, where ${}_{-A}X \in \{+, -, 0\}^E$ is the vector defined as follows:

$$
{}_{-A}X(e) := \begin{cases} -X(e) & \text{for } e \in A, \\ X(e) & \text{for } e \notin A. \end{cases}
$$

Definition 2.4 For $r, n \in \mathbb{N}$ $(n \geq r+1)$, the alternating matroid $A_{r,n}$ is an oriented matroid of rank r on a ground set $\{e_1, \dots, e_n\}$ with a chirotope χ such that $\chi(e_{i_1}, \dots, e_{i_r}) = +$ for all $1 \leq i_1 < i_2 < \cdots < i_r \leq n$.

Every alternating matroid is known to be convex and realizable. For more details on alternating matroids, see [2, Section 9.4].

2.2 Definitions on symmetry

2.2.1 Geometric symmetry of point configurations

Let $P := \{\boldsymbol{p}_1, \dots, \boldsymbol{p}_n\} \subseteq \mathbb{R}^d$ be a d-dimensional point configuration. A permutation σ on $\{1, \dots, n\}$ is a *geometric symmetry* of P if there exists an affine transformation f such that

$$
f(\boldsymbol{p}_i) = \boldsymbol{p}_{\sigma(i)} \text{ for } i = 1, \dots, n.
$$

Here, we present some other equivalent formulations. Let $\boldsymbol{v}_1, \dots, \boldsymbol{v}_n \in \mathbb{R}^{d+1}$ be the associated vector configuration of P. A permutation σ on $\{1, \dots, n\}$ is a geometric symmetry of P if and only if there exists a linear transformation A such that

$$
A\boldsymbol{v}_i = \boldsymbol{v}_{\sigma(i)} \text{ for } i = 1, \dots, n.
$$

Actually, this condition is equivalent to the condition that the following relations are satified:

$$
\det(\boldsymbol{v}_{\sigma(i_1)}, \dots, \boldsymbol{v}_{\sigma(i_{d+1})}) = \det(\boldsymbol{v}_{i_1}, \dots, \boldsymbol{v}_{i_{d+1}}) \text{ for all } 1 \leq i_1, \dots, i_r \leq n \cdots \text{ (G1), or}
$$

$$
\det(\boldsymbol{v}_{\sigma(i_1)}, \dots, \boldsymbol{v}_{\sigma(i_{d+1})}) = -\det(\boldsymbol{v}_{i_1}, \dots, \boldsymbol{v}_{i_{d+1}}) \text{ for all } 1 \leq i_1, \dots, i_r \leq n. \cdots \text{ (G2)}
$$

The permutation σ is called a *(geometric) rotational symmetry* of P if (G1) holds. On the other hand, σ is called a *(geometric) reflection symmetry* of P if (G2) holds.

2.2.2 Symmetry of oriented matroids

For an oriented matroid $\mathcal{M} = (\{1, \dots, n\}, \{\chi, -\chi\})$ of rank r, a permutation σ on E is a *symmetry* of $\mathcal{M}$ if $\mathcal{M}$ is invariant under σ, i.e., the following holds:

$$\chi(\sigma(i_1), \dots, \sigma(i_r)) = \chi(i_1, \dots, i_r) \text{ for all } 1 \leq i_1, \dots, i_r \leq n \cdots \text{ (O1), or}$$

$$\chi(\sigma(i_1), \dots, \sigma(i_r)) = -\chi(i_1, \dots, i_r) \text{ for all } 1 \leq i_1, \dots, i_r \leq n. \cdots \text{ (O2)}$$

The permutation σ is called a *(combinatorial) rotational symmetry* of $\mathcal{M}$ if Condition (O1) holds. We call σ a *(combinatorial) reflection symmetry* of $\mathcal{M}$ if it satisfies Condition (O2). The group formed by all rotational symmetries of $\mathcal{M}$ is called the *rotational symmetry group* of $\mathcal{M}$ and is denoted by $R(\mathcal{M})$. The group formed by all symmetries of $\mathcal{M}$ is called the symmetry group of $\mathcal{M}$ and is denoted by $G(\mathcal{M})$.

2.2.3 Combinatorial symmetry of point configurations

A permutation σ on $\{1, \dots, n\}$ is an *oriented-matroid symmetry (OM-symmetry)* of P if the oriented matroid associated to P is invariant under σ i.e.,

$$\text{sign}(\det(\boldsymbol{v}_{\sigma(i_1)}, \dots, \boldsymbol{v}_{\sigma(i_{d+1})})) = \text{sign}(\det(\boldsymbol{v}_{i_1}, \dots, \boldsymbol{v}_{i_{d+1}})) \text{ for all } 1 \leq i_1, \dots, i_r \leq n, \text{ or}$$

$$\text{sign}(\det(\boldsymbol{v}_{\sigma(i_1)}, \dots, \boldsymbol{v}_{\sigma(i_{d+1})})) = -\text{sign}(\det(\boldsymbol{v}_{i_1}, \dots, \boldsymbol{v}_{i_{d+1}})) \text{ for all } 1 \leq i_1, \dots, i_r \leq n.$$

On the other hand, σ is a *matroidal symmetry (M-symmetry)* if the matroid associated to P is invariant under σ, i.e.,

$$|\chi_P(\sigma(i_1), \dots, \sigma(i_{d+1}))| = |\chi_P(i_1, \dots, i_{d+1})| \text{ for all } 1 \leq i_1 < \dots < i_{d+1} \leq n.$$

OM-symmetries and M-symmetries are often called *combinatorial symmetries.*

It can easily be checked that every geometric symmetry induces a combinatorial symmetry. Given a combinatorial symmetry σ of P, σ is said to be *geometrically realizable* if there is a point configuration P' having the same oriented matroid (or matroid) as P and the geometric symmetry σ. In general, every combinatorial symmetry is not geometrically realizable as explained in the previous section. We will study this issue for matroids further in the following section.

3 A gap between matroidal symmetry and geometric symmetry of point configurations

3.1 A rank 4 matroid with 8 elements having an M-symmetry that cannot be realized geometrically

Let us consider a point configuration $P = \{\boldsymbol{p}_1, \boldsymbol{p}_2, \dots, \boldsymbol{p}_8\} \subseteq \mathbb{R}^3$ defined by

$$(\boldsymbol{p}_1, \boldsymbol{p}_2, \dots, \boldsymbol{p}_8) = \begin{pmatrix} 0 & 0 & 1 & 1 & 0 & 0 & \frac{1}{4} & \frac{1}{5} \\ 0 & 0 & 0 & 0 & 1 & 1 & \frac{1}{4} & \frac{1}{5} \\ 0 & 1 & 0 & 1 & 0 & 1 & -1 & 2 \end{pmatrix}.$$

The associated matroid M_P is described as follows.

```
1111211121121231112112123112123123411121121231121231234112123123412345
2223322332334442233233444233444555522332334442334445555233444555566666
3344434445555553444555555666666666634445555556666666666777777777777777
4555566666666667777777777777777777788888888888888888888888888888888888
0********0****0*******************************************************
```

Here, we are assigning $*$ (resp. 0) to row $ijkl$ if the set $\{i,j,k,l\}$ is a basis (resp. non-basis) of M_P. The associated matroid M_P has a symmetry σ:

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\ 3 & 4 & 5 & 6 & 1 & 2 & 7 & 8 \end{pmatrix}$$

In the following, we prove that M_P cannot be realized as a point configuration with a geometric symmetry σ. For $S \subseteq \mathbb{R}^3$, we will denote by $\mathrm{aff}(S)$ (resp. $\mathrm{conv}(S), \mathrm{relint}(S)$) the affine hull (resp. convex hull, relative interior) of S.

We assume that there is an affine automorphism f inducing σ. Then $C := \mathrm{conv}\{\boldsymbol{p}_1, \boldsymbol{p}_3, \boldsymbol{p}_5\}$ is setwise invariant under f. Note that f has a fixed point $\boldsymbol{w}_0$ in $\mathrm{relint}(\mathrm{conv}\{\boldsymbol{p}_1, \boldsymbol{p}_3, \boldsymbol{p}_5\})$ ($\boldsymbol{w}_0 = (\boldsymbol{p}_1 + \boldsymbol{p}_3 + \boldsymbol{p}_5)/3$). Since both $\boldsymbol{w}_0$ and $\boldsymbol{p}_8$ are invariant under f, $\mathrm{aff}(\{\boldsymbol{w}_0, \boldsymbol{p}_8\})$ is pointwise invariant under f. The same applies to $\mathrm{aff}(\{\boldsymbol{w}_0, \boldsymbol{p}_7\})$. Assume that $\boldsymbol{p}_7 \notin \mathrm{aff}(\{\boldsymbol{w}_0, \boldsymbol{p}_8\})$. Then $D := \mathrm{aff}(\{\boldsymbol{w}_0, \boldsymbol{p}_7, \boldsymbol{p}_8\})$ is 2-dimensional and pointwise invariant under f. Note that $\dim(\mathrm{aff}(C) \cap D) = 1$ since $\mathrm{aff}(C) \cap D \neq \phi$. Thus the restriction $f|_{\mathrm{aff}(C)}$ of f to $\mathrm{aff}(C)$ fixes a 1-dimensional space pointwisely. This contradicts to the fact that every non-trivial rational symmetry in the 2-dimensional Euclidean space has a unique fixed point. Therefore, we have $\boldsymbol{p}_7 \in \mathrm{aff}(\{\boldsymbol{w}_0, \boldsymbol{p}_8\})$.

Now we assume that lines $\mathrm{aff}(\{\boldsymbol{p}_1, \boldsymbol{p}_2\})$ and $\mathrm{aff}(\{\boldsymbol{p}_3, \boldsymbol{p}_4\})$ are parallel. Then lines $\mathrm{aff}(\{\boldsymbol{p}_1, \boldsymbol{p}_2\})$, $\mathrm{aff}(\{\boldsymbol{p}_3, \boldsymbol{p}_4\})$ and $\mathrm{aff}(\{\boldsymbol{p}_5, \boldsymbol{p}_6\})$ are all parallel. Therefore, lines $\mathrm{aff}(\{\boldsymbol{p}_1, \boldsymbol{p}_2\})$ and $\mathrm{aff}(\{\boldsymbol{w}_0, \boldsymbol{w}_1\})(= \mathrm{aff}(\{\boldsymbol{p}_7, \boldsymbol{p}_8\}))$ are also parallel and thus $\chi_P(1,2,7,8) = 0$, a contradiction.

Therefore, lines $\mathrm{aff}(\{\boldsymbol{p}_1, \boldsymbol{p}_2\})$ and $\mathrm{aff}(\{\boldsymbol{p}_3, \boldsymbol{p}_4\})$ are not parallel. Then lines $\mathrm{aff}(\{\boldsymbol{p}_1, \boldsymbol{p}_2\})$, $\mathrm{aff}(\{\boldsymbol{p}_3, \boldsymbol{p}_4\})$ and $\mathrm{aff}(\{\boldsymbol{p}_5, \boldsymbol{p}_6\})$ are all non-parallel. In this case, lines $\mathrm{aff}(\{\boldsymbol{p}_1, \boldsymbol{p}_2\})$ and $\mathrm{aff}(\{\boldsymbol{p}_3, \boldsymbol{p}_4\})$ intersect because points $\boldsymbol{p}_1$,$\boldsymbol{p}_2$,$\boldsymbol{p}_3$,$\boldsymbol{p}_4$ are on the same plane. The same apply to $\mathrm{aff}(\{\boldsymbol{p}_3, \boldsymbol{p}_4\})$ and $\mathrm{aff}(\{\boldsymbol{p}_5, \boldsymbol{p}_6\})$, and $\mathrm{aff}(\{\boldsymbol{p}_1, \boldsymbol{p}_2\})$ and $\mathrm{aff}(\{\boldsymbol{p}_5, \boldsymbol{p}_6\})$. Those three intersection points are in fact the same. This can be proved by the following simple computation.

$$\begin{aligned}
&\mathrm{aff}(\{\boldsymbol{p}_1, \boldsymbol{p}_2\}) \cap \mathrm{aff}(\{\boldsymbol{p}_3, \boldsymbol{p}_4\}) \\
&= (\mathrm{aff}(\{\boldsymbol{p}_1, \boldsymbol{p}_2, \boldsymbol{p}_3, \boldsymbol{p}_4\}) \cap \mathrm{aff}(\{\boldsymbol{p}_1, \boldsymbol{p}_2, \boldsymbol{p}_5, \boldsymbol{p}_6\})) \cap (\mathrm{aff}(\{\boldsymbol{p}_1, \boldsymbol{p}_2, \boldsymbol{p}_3, \boldsymbol{p}_4\}) \cap \mathrm{aff}(\{\boldsymbol{p}_3, \boldsymbol{p}_4, \boldsymbol{p}_5, \boldsymbol{p}_6\})) \\
&= \mathrm{aff}(\{\boldsymbol{p}_1, \boldsymbol{p}_2, \boldsymbol{p}_3, \boldsymbol{p}_4\}) \cap \mathrm{aff}(\{\boldsymbol{p}_3, \boldsymbol{p}_4, \boldsymbol{p}_5, \boldsymbol{p}_6\}) \cap \mathrm{aff}(\{\boldsymbol{p}_1, \boldsymbol{p}_2, \boldsymbol{p}_5, \boldsymbol{p}_6\}).
\end{aligned}$$

This intersection is invariant under A and thus is on $\mathrm{aff}(\{\boldsymbol{p}_7, \boldsymbol{p}_8\})$.
From the above discussion, lines $\mathrm{aff}(\{\boldsymbol{p}_5, \boldsymbol{p}_6\})$ and $\mathrm{aff}(\{\boldsymbol{p}_7, \boldsymbol{p}_8\})$ have the intersection and thus $\boldsymbol{p}_5$,$\boldsymbol{p}_6$, $\boldsymbol{p}_7$,$\boldsymbol{p}_8$ are on the same plane. This leads $\chi_P(5,6,7,8) = 0$, a contradiction.



Figure 1: Case1 [left], Case2 [right]

3.2 General construction

In a similar way to the proof in the previous subsection, it can be proved that from an arbitrary 2-dimensional configuration of n points with a non-trivial geometric rotational symmetry of order $m > 2$, we can construct a 3-dimensional configuration of $2n+2$ points with an M-symmetry that cannot be realized geometrically.

Theorem 3.1 Let $P := \{\boldsymbol{p}_1, \ldots, \boldsymbol{p}_n\} \subseteq \mathbb{R}^2$ be a point configuration with a non-trivial geometric rotational symmetry σ of order $m > 2$. Then let us consider a point configuration $Q := \{\boldsymbol{q}_1, \ldots, \boldsymbol{q}_{2n+2}\} \subseteq \mathbb{R}^3$ where $\boldsymbol{q}_i := \begin{pmatrix} \boldsymbol{p}_i \\ 0 \end{pmatrix}, \boldsymbol{q}_{n+i} := \begin{pmatrix} \boldsymbol{p}_i \\ 1 \end{pmatrix}$ for $i = 1, \ldots, n$ and $\boldsymbol{q}_{2n+1}, \boldsymbol{q}_{2n+2}$ are *generic* points such that the line $\operatorname{aff}(\{\boldsymbol{q}_{2n+1}, \boldsymbol{q}_{2n+2}\})$ is not parallel to $\operatorname{aff}(\{\boldsymbol{p}_1, \boldsymbol{p}_{n+1}\})$. Here we say a point $\boldsymbol{r}$ in a point configuration R is generic if $\boldsymbol{r} \notin \operatorname{aff}(\{\boldsymbol{r}_1, \boldsymbol{r}_2\})$ for all $\boldsymbol{r}_1, \boldsymbol{r}_2 \in R \setminus \{\boldsymbol{r}\}$. Then Q has a matroidal symmetry that cannot be realized geometrically.

PROOF: Let M_Q is the matroid of Q. First of all, note that M_Q has a symmetry

$$\tau := \begin{pmatrix} 1 & \ldots & n & n+1 & \ldots & 2n & 2n+1 & 2n+2 \\ \sigma(1) & \ldots & \sigma(n) & n+\sigma(1) & \ldots & n+\sigma(n) & 2n+1 & 2n+2 \end{pmatrix}.$$

Suppose that there is a realization $R = (\boldsymbol{r}_1, \ldots, \boldsymbol{r}_{2n+2}) \in \mathbb{R}^{3\times(2n+2)}$of M_Q and an affine transformation f of $\mathbb{R}^3$ with $f(\boldsymbol{r}_i) = \boldsymbol{r}_{\tau(i)}$ for $i = 1, \ldots, 2n+2$.
Let $\boldsymbol{r}_{i_0}$ be an extreme point of $C_1 := \operatorname{conv}\{\boldsymbol{r}_1, \ldots, \boldsymbol{r}_n\}$. Then $\boldsymbol{r}_{i_0+n}$ is also an extreme point of $C_2 := \operatorname{conv}\{\boldsymbol{r}_{n+1}, \ldots, \boldsymbol{r}_{2n}\}$. Consider the orbits $O_1 = \{\boldsymbol{r}_{i_0}, \ldots, \boldsymbol{r}_{i_m}\}$ of $\boldsymbol{r}_{i_0}$ and $O_2 = \{\boldsymbol{r}_{i_0+n}, \ldots, \boldsymbol{r}_{i_m+n}\}$ of $\boldsymbol{r}_{i_0+n}$. Notice that $|O_1| = |O_2| \geq 3$.

Now let us consider $\boldsymbol{w}_0 := \frac{1}{m+1}(\boldsymbol{r}_{i_0} + \cdots + \boldsymbol{r}_{i_m})$ and $\boldsymbol{w}_1 := \frac{1}{m+1}(\boldsymbol{r}_{i_0+n} + \cdots + \boldsymbol{r}_{i_m+n})$. Since $\boldsymbol{w}_0$ and $\boldsymbol{w}_1$ are invariant under f, the line $\operatorname{aff}(\{\boldsymbol{w}_0, \boldsymbol{r}_{2n+1}\})$ is pointwise invariant under f. Note that $\boldsymbol{r}_{2n} \in \operatorname{aff}(\{\boldsymbol{w}_0, \boldsymbol{r}_{2n+1}\})$. Otherwise, we have a 2-dimensional space $D := \operatorname{aff}(\{\boldsymbol{w}_0, \boldsymbol{r}_{2n}, \boldsymbol{r}_{2n+1}\})$ pointwise invariant under f. Since $D \cap \operatorname{aff}(C_1) \neq \phi$, it holds that $\dim(\operatorname{aff}(C_1) \cap D) = 1$. Thus the restriction $f|_{\operatorname{aff}(C_1)}$ of f to $\operatorname{aff}(C_1)$ fixes a 1-dimensional space pointwisely, which is a contradiction. Therefore, it holds that $\boldsymbol{r}_{2n} \in \operatorname{aff}(\{\boldsymbol{w}_0, \boldsymbol{r}_{2n+1}\})$. Similarly, we have $\boldsymbol{w}_1 \in \operatorname{aff}(\{\boldsymbol{w}_0, \boldsymbol{r}_{2n+1}\})$.

Now assume that lines $\operatorname{aff}(\{\boldsymbol{r}_{i_0}, \boldsymbol{r}_{i_0+n}\})$ and $\operatorname{aff}(\{\boldsymbol{r}_{i_1}, \boldsymbol{r}_{i_1+n}\})$ are parallel. Then lines $\operatorname{aff}(\{\boldsymbol{r}_{i_0}, \boldsymbol{r}_{i_0+n}\})$, $\operatorname{aff}(\{\boldsymbol{r}_{i_m}, \boldsymbol{r}_{i_m+n}\})$ are all parallel. This implies that lines $\operatorname{aff}(\{\boldsymbol{r}_{i_0}, \boldsymbol{r}_{i_0+n}\})$ and $\operatorname{aff}(\{\boldsymbol{w}_0, \boldsymbol{w}_1\})$ are parallel and thus $\chi_P(i_0, i_0+n, 2n+1, 2n+2) = 0$, a contradiction.

Let us consider the case when lines $\operatorname{aff}(\{\boldsymbol{r}_{i_0}, \boldsymbol{r}_{i_0+n}\})$ and $\operatorname{aff}(\{\boldsymbol{r}_{i_1}, \boldsymbol{r}_{i_1+n}\})$ are not parallel. Then lines $\operatorname{aff}(\{\boldsymbol{r}_{i_0}, \boldsymbol{r}_{i_0+n}\})$ and $\operatorname{aff}(\{\boldsymbol{r}_{i_1}, \boldsymbol{r}_{i_1+n}\})$ have an intersection point since $\chi_P(i_0, i_0+n, i_1, i_1+n) = 0$. The same applies to $\operatorname{aff}(\{\boldsymbol{r}_{i_1}, \boldsymbol{r}_{i_1+n}\})$ and $\operatorname{aff}(\{\boldsymbol{r}_{i_2}, \boldsymbol{r}_{i_2+n}\})$, and $\operatorname{aff}(\{\boldsymbol{r}_{i_0}, \boldsymbol{r}_{i_0+n}\})$ and $\operatorname{aff}(\{\boldsymbol{r}_{i_2}, \boldsymbol{r}_{i_2+n}\})$. These three intersection points are in fact the same. This follows from the following relation:

$$\begin{aligned}
&\operatorname{aff}(\{\boldsymbol{r}_{i_0}, \boldsymbol{r}_{i_0+n}\}) \cap \operatorname{aff}(\{\boldsymbol{r}_{i_1}, \boldsymbol{r}_{i_1+n}\}) \\
&= \operatorname{aff}(\{\boldsymbol{r}_{i_0}, \boldsymbol{r}_{i_0+n}, \boldsymbol{r}_{i_1}, \boldsymbol{r}_{i_1+n}\}) \cap \operatorname{aff}(\{\boldsymbol{r}_{i_1}, \boldsymbol{r}_{i_1+n}, \boldsymbol{r}_{i_2}, \boldsymbol{r}_{i_2+n}\}) \cap \operatorname{aff}(\{\boldsymbol{r}_{i_0}, \boldsymbol{r}_{i_0+n}, \boldsymbol{r}_{i_2}, \boldsymbol{r}_{i_2+n}\}) \\
&= \operatorname{aff}(\{\boldsymbol{r}_{i_1}, \boldsymbol{r}_{i_1+n}\}) \cap \operatorname{aff}(\{\boldsymbol{r}_{i_2}, \boldsymbol{r}_{i_2+n}\}) \\
&= \operatorname{aff}(\{\boldsymbol{r}_{i_0}, \boldsymbol{r}_{i_0+n}\}) \cap \operatorname{aff}(\{\boldsymbol{r}_{i_2}, \boldsymbol{r}_{i_2+n}\})
\end{aligned}$$

Repeating the same argument, we notice that

$$\operatorname{aff}(\{\boldsymbol{r}_{i_0}, \boldsymbol{r}_{i_0+n}\}) \cap \cdots \cap \operatorname{aff}(\{\boldsymbol{r}_{i_m}, \boldsymbol{r}_{i_m+n}\})$$

is non-empty. Since it is a fixed point of f, it is on the line $\operatorname{aff}(\{\boldsymbol{w}_0, \boldsymbol{w}_1\}) = \operatorname{aff}(\{\boldsymbol{r}_{2n+1}, \boldsymbol{r}_{2n+2}\})$. This implies $\chi_P(i_0, i_0+n, 2n+1, 2n+2) = 0$, a contradiction. $\square$

Corollary 3.2 For $n \geq 3$, from any 2-dimensional configuration of n points with a geometric rotational symmetry σ, we can construct a 3-dimensional configuration of $2n+2$ points with an M-symmetry that cannot be realized geometrically.

4 Fixed point theorems for rotational symmetries of oriented matroids

The construction of a gap between M-symmetry and geometric symmetry in the previous section is based on the fact that every non-trivial rotation in the 2-dimensional Euclidean space has a unique fixed point, but matroids do not have the corresponding property. A natural question is whether oriented matroids have the corresponding property or not. In this section, we study the corresponding property for oriented matroids.

4.1 Uniqueness property

The following is an oriented-matroid analogue of the uniqueness of a fixed point of a non-trivial rotational symmetry in the 2-dimensional Euclidean space.

Proposition 4.1 Let $\mathcal{M} = (\{1,\dots,n\},\{\chi,-\chi\})$ be a loopless oriented matroid of rank 3 with a rotational symmetry σ of order a. Assume that σ is a *non-trivial rotational symmetry*, i.e., there exists $x \in \{1,\dots,n\}$ such that x and $\sigma(x)$ are not parallel. Then, if $\sigma(p)=p, \sigma(q)=q$, it holds that p and q are parallel or antiparallel, i.e., $\chi(p,q,x)=0$ for all $x \in \{1,\dots,n\}$.

Proof:
Assume that p and q are neither parallel nor antiparallel. Then, we have $rank(Flat(\{p,q\}))=2$. First, let us consider $x \in \{1,\dots,n\}$ such that $\chi(p,q,x) \neq 0$. For any $s,t \in \mathbb{N}$, the following holds:

$$\{\chi(p,q,x)\chi(p,\sigma^s(x),\sigma^{s+t}(x)), -\chi(p,q,\sigma^s(x))\chi(p,x,\sigma^{s+t}(x)), \chi(p,q,\sigma^{s+t}(x))\chi(p,x,\sigma^s(x))\} \\ \supseteq \{+,-\} \text{ or } =\{0\}$$

by the chirotope axiom. Therefore,

$$\{\chi(p,q,x)\chi(p,x,\sigma^t(x)), -\chi(p,q,x)\chi(p,x,\sigma^{s+t}(x)), \chi(p,q,x)\chi(p,x,\sigma^s(x))\} \\ \supseteq \{+,-\} \text{ or } =\{0\}.$$

Since $\chi(p,q,x) \neq 0$,

$$\{\chi(p,x,\sigma^t(x)), -\chi(p,x,\sigma^{s+t}(x)), \chi(p,x,\sigma^s(x))\} \supseteq \{+,-\} \text{ or } =\{0\}.$$

Therefore, for any s,t such that $\chi(p,x,\sigma^t(x)) = \chi(p,x,\sigma^s(x))$, we have

$$\chi(p,x,\sigma^s(x)) = \chi(p,x,\sigma^{s+t}(x)).$$

By induction, we have

$$\chi(p,x,\sigma(x)) = \chi(p,x,\sigma^l(x)) \text{ for all } l \geq 1.$$

Since $\chi(p,x,\sigma^a(x)) = \chi(p,x,x) = 0$ and thus

$$\chi(p,x,\sigma^l(x)) = 0 \text{ for all } l \geq 0.$$

Similarly, it holds that

$$\chi(q,x,\sigma^l(x)) = 0 \text{ for all } l \geq 0.$$

By the chirotope axioms,

$$\{\chi(x,y,\sigma(x))\chi(x,p,q), -\chi(x,y,p)\chi(x,\sigma(x),q), \chi(x,y,q)\chi(x,\sigma(x),p)\} \\ = \{0, \chi(x,y,\sigma(x))\chi(x,p,q)\} \supseteq \{+,-\} \text{ or } =\{0\}$$

for any $y \in E$. Since $\chi(p,q,x) \neq 0$, it holds that $\chi(x,y,\sigma(x)) = 0$ for any $y \in \{1,\dots,n\}$. Therefore, x and $\sigma(x)$ are parallel or antiparallel. In fact, it cannot occur that x and $\sigma(x)$ are antiparallel since σ is a rotational symmetry.

Next, let us consider $x \in \{1,\dots,n\}$ such that $\chi(p,q,x) = 0$. Assume that x and $\sigma(x)$ are not parallel (and are not antiparallel). Then there exists $y \in \{1,\dots,n\}$ such that $\chi(y,x,\sigma(x)) \neq 0$. Note that $x, \sigma(x) \in \mathit{Flat}(\{p,q\})$. Since $y \notin \mathit{Flat}(\{p,q\})$, it turns out that y and $\sigma(y)$ are parallel by applying the same argument as the above. For any $s,t \in \mathbb{N}$, the following holds:

$$\begin{aligned}\{\chi(y,p,x)\chi(y,\sigma^s(x),\sigma^{s+t}(x)), -\chi(y,p,\sigma^s(x))\chi(y,x,\sigma^{s+t}(x)), \chi(y,p,\sigma^{s+t}(x))\chi(y,x,\sigma^s(x))\}\\ \supseteq \{+,-\} \text{ or } = \{0\}\end{aligned}$$

by the chirotope axiom. Note that $\chi(y,p,x) = \chi(y,p,\sigma^s(x)) = \chi(y,p,\sigma^{s+t}(x))$. If $\chi(p,x,y) \neq 0$, then $\chi(y,x,\sigma(x)) = 0$ similarly to the above discussion, a contradiction. Therefore, $\chi(p,x,y) = 0$. Since p and x are not parallel nor antiparallel, there exists $z \in \{1,\dots,n\}$ such that $\chi(p,x,z) \neq 0$. However, it holds that

$$\begin{aligned}&\{\chi(p,q,y)\chi(p,x,z), -\chi(p,q,x)\chi(p,y,z), \chi(p,q,z)\chi(p,y,x)\}\\ &= \{\chi(p,q,y)\chi(p,x,z)(\neq 0), 0\} \supseteq \{+,-\} \text{ or } = \{0\},\end{aligned}$$

a contradiction. Therefore, x and $\sigma(x)$ are parallel.

As a conclusion, σ is a trivial rotational symmetry, which is a contradiction. Therefore, p and q are parallel or antiparallel. □

The above proof can easily be extended to the following theorem.

Theorem 4.2 *Let $\mathcal{M} = (\{1,\dots,n\}, \{\chi,-\chi\})$ be a loopless oriented matroid of rank r with a non-trivial rotational symmetry σ. Let us set $\mathit{Fix}(\sigma) := \{e \in \{1,\dots,n\} \mid \sigma(e) = e\}$. Then, we have $\mathit{rank}(\mathcal{M}|_{\mathit{Fix}(\sigma)}) \leq r-2$.*

One of useful applications of the above theorem is as follows.

Corollary 4.3 *Let $\mathcal{M}$ be a loopless oriented matroid of rank r. For $\sigma,\tau \in R(\mathcal{M})$, if $\sigma|_X = \tau|_X$ for $X \subseteq E$ with $\mathit{rank}(\mathcal{M}|_X) \geq r-1$, then $\sigma = \tau$.*

PROOF: Assume that $\sigma|_X = \tau|_X$ but $\sigma \neq \tau$. Then $\sigma\tau^{-1}$ is a non-trivial rotational symmetry of $\mathcal{M}$. However, $\sigma\tau^{-1}$ fixes X pointwisely. This is a contradiction. □

Remark 4.4 Uniqueness in the the following sense does not hold.

> Let $\mathcal{M}$ be a rank 3 oriented matroid on a ground set $\{1,\dots,n\}$ with a non-trivial rotational symmetry σ. If $\mathcal{M}\cup p$ and $\mathcal{M}\cup q$ have rotational symmetries $\sigma' := \begin{pmatrix} 1 & 2 & \cdots & n & p \\ \sigma(1) & \sigma(2) & \cdots & \sigma(n) & p \end{pmatrix}$ and $\sigma'' := \begin{pmatrix} 1 & 2 & \cdots & n & q \\ \sigma(1) & \sigma(2) & \cdots & \sigma(n) & q \end{pmatrix}$ respectively, then we have $\mathcal{M}\cup p = \mathcal{M}\cup q$.

A counterexample is given in Figure 2.

4.2 Existence property

Now we prove existence of a fixed point of a rotational symmetry of a matroid polytope. This part is not proved for general oriented matroids, but it is sufficient for the remaining part of this paper.



Figure 2: Fixed points for oriented matroids

Theorem 4.5 Let $\mathcal{M} = (\{1, \ldots, n\}, \{\chi, -\chi\})$ be a simple acyclic oriented matroid with all elements extreme points. Then there exists a single element extension $\mathcal{M} \cup \{n+1\} = (\{1, \ldots, n+1\}, \{\chi', -\chi'\})$ of $\mathcal{M}$ which is invariant under $\sigma' := \begin{pmatrix} 1 & 2 & \cdots & n & n+1 \\ \sigma(1) & \sigma(2) & \cdots & \sigma(n) & n+1 \end{pmatrix}$ for all $\sigma \in R(\mathcal{M})$.

Proof: A single element extensions of an oriented matroid is determined by cocircuit signatures and there is a good characterization of cocircuit signatures that define a single element extension [9]. The reader can find a survey on this topic in [2, Section 7]. In this proof, we construct a desired single element extension of $\mathcal{M}$ using the characterization by Las Vergnas [9].

Let $\mathcal{C}^*$ be the set of cocircuits of $\mathcal{M}$. We define function $n_+ : \mathcal{C}^* \to \mathbb{N}$ and $n_- : \mathcal{C}^* \to \mathbb{N}$ by

$$n_+(c) = |\{i \in \{1, \ldots, n\} \mid c(i) = +\}|,$$

$$n_-(c) = |\{i \in \{1, \ldots, n\} \mid c(i) = -\}|$$

for $c \in \mathcal{C}^*$, and define cocircuit signatures $l : \mathcal{C}^* \to \{+, -, 0\}$ as follows.

$$l(c) = \begin{cases} + & \text{if } n_+(c) > n_-(c), \\ - & \text{if } n_+(c) < n_-(c), \\ 0 & \text{if } n_+(c) = n_-(c). \end{cases}$$

We prove that l is a localization. Let L be a coline of $\mathcal{M}$ and $c_1 c_2 \ldots c_m$ be a cocircuit (on L) ordering along L. Without loss of generality, we assume that c_1, c_m are non-negative cocircuits (recall that $\mathcal{M}$ is convex). Then, we have

$$n_+(c_1), n_+(c_m) \geq n_+(c_2), \ldots, n_+(c_{m-1}).$$

Then, $2 \leq^{\exists} j \leq n$ s.t.

$$n_+(c_1) > n_+(c_2) > \cdots > n_+(c_j) = n_+(c_{j+1}) = 0 < \cdots < n_+(c_m) = n_+(c_1) \text{ and}$$

$$n_-(c_1) < n_-(c_2) < \cdots < n_-(c_j) = n_-(c_{j+1}) = n > \cdots > n_-(c_m) = n_-(c_1).$$

This is proved as follows. Since $c_1 c_2 \ldots c_m$ is a cocircuit ordering along a coline, sign pattern $c_1(e) c_2(e) \ldots c_m(e)$ is one of the following types:

$$0 - - \cdots - 0 + + \cdots + \quad \text{or} \quad + + \cdots + 0 - - \cdots - 0 + + \cdots + \quad \text{or} \quad + + \cdots + 0 - - \cdots - 0$$

for all $1 \leq e \leq n$, where $c_1(e), \ldots, c_m(e)$ denote the e-th component of $c_1, \ldots, c_m$. In any cases, there are exactly two k s.t. $c_k(e) = 0$. Let c_j be a non-positive cocircuit (note that $n(c_j) = 0$). It can easily seen that if $c_{k_1}(e) = c_{k_2}(e) = 0$ for $k_1 < k_2$ then $1 \leq k_1 \leq j$ and $j \leq k_2 \leq n$. Therefore, the sign sequence $c_1(e)c_2(e)...c_j(e)$ is decreasing and the sign sequence $c_j(e)c_{j+1}(e)...c_m(e)$ is increasing for all $1 \leq e \leq n$. This proves the above claim.

From the above relations, we have

$$\begin{aligned} & n_+(c_1) - n_-(c_1) > n_+(c_2) - n_-(c_2) > \cdots > n_+(c_j) - n_-(c_j) = n_+(c_{j+1}) - n_-(c_{j+1}) = -n \\ & < \cdots < n_+(c_m) - n_-(c_m). \end{aligned}$$

Therefore, there exist j_1, j_2 such that

$$\begin{cases} n_+(c_i) > n_-(c_i) & \text{for } 0 \leq i < j_1,\ j_2 \leq i \leq m, \\ n_+(c_i) < n_-(c_i) & \text{for } j_1 \leq i < j_2 \end{cases}$$

or

$$\begin{cases} n_+(c_i) > n_-(c_i) & \text{for } 0 \leq i < j_1,\ j_2 + 1 \leq i \leq m, \\ n_+(c_i) < n_-(c_i) & \text{for } j_1 + 1 \leq i < j_2, \\ n_+(c_i) = n_-(c_i) & \text{for } i = j_1, j_2. \end{cases}$$

Additionally, we have $l(-c) = -l(c)$ for all $c \in \mathcal{C}^*$. This implies that l certainly defines a single element extension of $\mathcal{M}$. Let us denote the single element extension by $\mathcal{M} \cup \{n+1\}$. Clearly, the oriented matroid $\mathcal{M} \cup \{n+1\}$ is invariant under σ'. $\Box$

Remark 4.6 The oriented matroid $\mathcal{M}'$ in the above theorem is also simple and acyclic. If $\mathcal{M}$ is cosimple, $\mathcal{M}'$ is also cosimple. However, $\mathcal{M}'$ is not a matroid polytope.

Remark 4.7 For a rank r matroid polytope with the non-trivial rotational symmetry group $R(\mathcal{M})$, we can take a fixed point q by the above theorem and can construct a rank $r-1$ oriented matroid $(\mathcal{M} \cup q)/q$, which is also invariant under $R(\mathcal{M})$. Let $\sigma \in R(\mathcal{M})$ be a rotational symmetry of $\mathcal{M}$ and χ a chirotope of $\mathcal{M}$. Then, the map $\chi' : E^{r-1} \to \{+, -, 0\}$ defined by

$$\chi'(i_1, \ldots, i_{r-1}) := \chi(q, i_1, \ldots, i_{r-1}) \text{ for } i_1, \ldots, i_{r-1} \in E$$

is a chirotope of $(\mathcal{M} \cup q)/q$. Then, it holds that

$$\sigma \cdot \chi'(i_1, \ldots, i_{r-1}) = \chi'(\sigma(i_1), \ldots, \sigma(i_{r-1})) = \chi(q, \sigma(i_1), \ldots, \sigma(i_{r-1})) = \chi(q, i_1, \ldots, i_{r-1}) = \chi'(i_1, \ldots, i_{r-1})$$

for all $i_1, \ldots, i_{r-1} \in E$. Therefore, σ is a rotational symmetry of $(\mathcal{M} \cup q)/q$. In this way, we can decrease the rank of an oriented matroid by one if it is convex, preserving rotational symmetries.

Remark 4.8 Consider a simple non-acyclic oriented matroid $\mathcal{M}$ of rank r on a ground set E where $R(\mathcal{M})$ acts on E transitively. Then, the oriented matroid $\mathcal{M}$ is totally cyclic and thus the dual oriented matroid $\mathcal{M}^*$ is acyclic. Since $R(\mathcal{M}^*)(= R(\mathcal{M}))$ acts on E transitively, $\mathcal{M}^*$ is a matroid polytope of rank $|E| - r$. Then, let us take a fixed point q and consider a single element extension $\mathcal{M}^* \cup q$. Note that $R(\mathcal{M}^*) = R((\mathcal{M}^* \cup q)/q)$ holds and that the contraction $(\mathcal{M}^* \cup q)/q$ is cyclic and thus totally cyclic by transitivity. This leads that the dual oriented matroid $\mathcal{N} := ((\mathcal{M}^* \cup q)/q)^*$ is acyclic. $\mathcal{N}$ is a rank $r+1$ matroid polytope by transitivity. Note that $\mathcal{N}$ is simple.

The above remarks show that classification of (rotational) symmetry groups of non-acyclic oriented matroids of rank r is closely related to that of acyclic oriented matroids (or matroid polytopes, equivalently) of rank $r+1$.

5 Symmetry groups of simple oriented matroids of rank 3 in a certain class

In this section, we investigate rotational symmetry groups of rank 3 oriented matroids. Results in this part will be used to investigate the rank 4 case in the next section.

Before investigating simple oriented matroids of rank 3, we briefly touch upon the rank 2 case. Every simple oriented matroid $\mathcal{M}$ of rank 2 on a ground set E can be represented as a (unsigned) point configuration on a 1-sphere. By the simple consideration, all the symmetries of $\mathcal{M}$ can be viewed as symmetries of the $|E|$-cycle graph. It is straightforward that $R(\mathcal{M})$ is a cyclic group and $G(\mathcal{M})$ is a dihedral group. We note here that a rank 2 simple acyclic oriented matroid cannot have a rotational symmetry of order $p > 2$.

Now let us move to the rank 3 case. The acyclic case is easy. Let $\mathcal{M}$ be a simple acyclic oriented matroid of rank 3 on a ground set E on which $R(\mathcal{M})$ acts transitively. Then, $\mathcal{M}$ is a rank 3 matroid polytope and thus a relabeling of the alternating matroid $A_{3,|E|}$. Therefore, the rotational symmetry group $R(\mathcal{M})$ is a cyclic group of order $|E|$. This leads that rotational symmetry groups of simple acyclic oriented matroids of rank 3 are covered by cyclic groups C_n $(n \geq 1)$.

To classify rotational symmetry groups of simple acyclic oriented matroids of rank 4, it is also necessary to investigate rotational symmetry groups of non-acyclic oriented matroids in certain classes. In the remaining part of this section, we will investigate some non-acyclic cases.

We will exclude the possibility that the cyclic group generated by a rotational symmetry of order $p > 2$ generates a rank 2 orbit. Note that such a situation cannot occur in the acyclic case. Let us call a rotational symmetry of a rank 3 oriented matroid a *proper rotational symmetry* if its order is 2 or it generates a rank 3 orbit.

5.1 Structures of rank 3 orbits under the actions of cyclic groups generated by proper rotational symmetries

Cyclic groups are groups of the simplest type. First, we investigate structures of orbits under the actions of cyclic groups generated by proper rotational symmetries.

Proposition 5.1 Let $\mathcal{M}$ be a simple (not necessarily acyclic) oriented matroid of rank 3 on a ground set E with a non-trivial rotational symmetry σ. If the cyclic group G generated by σ acts on E transitively, $\mathcal{M}$ is a relabeling of the alternating matroid $A_{3,|E|}$.

PROOF: It is sufficient to prove that $\mathcal{M}$ is a matroid polytope. Since the cyclic group G acts on E transitively, E can be written as $E = \{x, \sigma(x), \sigma^2(x), \ldots, \sigma^{n-1}(x)\}$, where $n := |E|$. Let us simply write i to describe $\sigma^i(x) \in E$ when there is no confusion.

> **Lemma:** Let $\mathcal{N}$ be a rank 3 simple oriented matroid on a ground set $F = \{1, \ldots, m\}$ $(m \geq 5)$ such that $\mathcal{N}|_{F-\{1\}}$ and $\mathcal{N}|_{F-\{m\}}$ are rank 3 matroid polytopes. Then $\mathcal{N}$ is acyclic.

Before proving the above lemma, let us note that $\mathcal{M}|_{E_4}$ is acyclic and thus a matroid polytope by transitivity. Let χ be a chirotope of $\mathcal{M}|_{E_4}$. Then, we have $\chi(1,2,3) = \chi(2,3,4)$. Simple case analysis on the values of $\chi(1,2,4)$ and $\chi(1,3,4)$ shows the existence of the positive covector $(++++)$.
(Proof of the above lemma) Assume that $\mathcal{N}$ is cyclic i.e., there exists an integer $1 \leq k \leq m$ and a vector X such that

$$\begin{cases} X(i_1) = X(i_2) = \cdots = X(i_k) = +, \\ X(a) = 0 \text{ for all } a \in F \setminus \{i_1, i_2, \ldots, i_k\}, \end{cases}$$

where $i_1, \ldots, i_k$ are numbers between 0 and $m+1$. If $i_1, \ldots, i_k > 1$, $X|_{F-\{1\}}$ is a vector of $M|_{F-\{1\}}$, a contradiction. This leads $1 \in \{i_1, \ldots, i_k\}$. For distinct numbers s, t, u between 1 and m, there exist a vector Y of $\mathcal{M}$ such that

$$\begin{cases} Y(1) = -, Y(s) = -, Y(t) = +, Y(u) = +, \\ Y(a) = 0 \text{ for all } a \in F \setminus \{1, s, t, u\}, \end{cases}$$

since $1, s, t, u$ are extreme points of the uniform oriented matroid $\mathcal{M}|_{F-\{m\}}$. By applying vector elimination to X, Y and 1, we obtain a vector Z of $\mathcal{M}$ such that

$$\begin{cases} Z(1) = 0, Z(s) \in \{+, -, 0\}, \\ Z(a) \in \{+, 0\} \text{ for all } a \in F \setminus \{1, s\}. \end{cases}$$

$Z|_{F\setminus\{1\}}$ is a vector of $\mathcal{M}|_{F\setminus\{1\}}$. This contradicts to the assumption that $\mathcal{M}|_{F-\{1\}}$ is a matroid polytope. This completes the proof of the lemma.

Now we set $E_m := \{1, 2, \dots, m\}$ for $m = 2, 3, \dots, n$. We prove that if $\mathcal{M}|_{E_m}$ is a matroid polytope, then $\mathcal{M}|_{E_{m+1}}$ is a matroid polytope, for $m = 4, \dots, n-1$.

By the above lemma, the oriented matroid $\mathcal{M}|_{E_{m+1}}$ is acyclic (note that $\mathcal{M}|_{E_{m+1}\setminus\{1\}}$ is a matroid polytope by transitivity). Assume that $\mathcal{M}|_{E_{m+1}}$ is not a matroid polytope.

(i) There exists a vector X of $\mathcal{M}|_{E_{m+1}}$ such that $X(m+1) = -$ and $X(a) = +$ for $i = 1, \dots, m$.
In this case, there exists a vector Y of $\mathcal{M}$ such that

$$\begin{cases} Y(m+1) = -, \\ Y(a) = + \text{ for all } a \in E_m, \\ Y(b) = 0 \text{ for all } E \setminus E_{m+1}. \end{cases}$$

Let us consider the vector $Y_1 := \sigma(Y)$, which satisfies

$$\begin{cases} Y_1(m+2) = -, \\ Y_1(a+1) = + \text{ for all } a \in E_m, \\ Y_1(b) = 0 \text{ for all } (E \setminus E_{m+2}) \cup \{1\}. \end{cases}$$

By applying vector elimination to Y, Y_1 and $m+1$, we obtain a vector Z_1 of $\mathcal{M}$ such that

$$\begin{cases} Z_1(m+2) = -, \\ Z_1(a) = + \text{ for all } a \in E_m, \\ Z_1(b) = 0 \text{ for all } (E \setminus E_m) \setminus \{m+2\}. \end{cases}$$

Let us consider the vector $Y_1 := \sigma^2(Y)$, which satisfies

$$\begin{cases} Y_2(m+3) = -, \\ Y_2(a+2) = + \text{ for all } a \in E_{m+2} \setminus \{1, 2\}, \\ Y_2(b) = 0 \text{ for all } (E \setminus E_{m+3}) \cup \{1, 2\}. \end{cases}$$

Let us take a vector W obtained by applying vector elimination to Y_2, Z_1 and $m+2$, and then a vector Z_2 obtained by applying vector elimination to W, Y and $m+1$.

$$\begin{cases} Z_2(m+3) = -, \\ Z_2(a) = + \text{ for all } a \in E_m, \\ Z_2(b) = 0 \text{ for all } (E \setminus E_m) \setminus \{m+3\}. \end{cases}$$

By repeating the same argument, we obtain a vector Z' of $\mathcal{M}$ such that

$$\begin{cases} Z'(n) = -, \\ Z'(a) = + \text{ for all } a \in E_m, \\ Z'(a) = 0 \text{ for all } a \in (E \setminus E_m) \setminus \{n\}. \end{cases}$$

The vector $Z'' := \sigma(Z')$ satisfies

$$\begin{cases} Z''(1) = -, \\ Z''(a) = + \text{ for } a = 2, \dots, m+1, \\ Z''(b) = 0 \text{ for } b = m+2, \dots, n. \end{cases}$$

By applying vector elimination to Y, Z'' and $m+1$, we obtain a vector U such that

$$\begin{cases} U(1) \in \{-, 0, +\}, \\ U(a) = + \text{ for } a = 2, \dots, m, \\ U(b) = 0 \text{ for } b = m+1, \dots, n. \end{cases}$$

This contradicts to the assumption that $\mathcal{M}|_{E_m}$ is a matroid polytope.

(ii) There exists a vector X of $\mathcal{M}|_{E_{m+1}}$ such that $X(k) = -$ and $X(a) = +$ for $E_{m+1} \setminus \{k\}$ $(2 \le k \le m+1)$. One can apply a similar argument to (i), keeping in mind that $\mathcal{M}|_{E_{m+1}}$ is acyclic. □

5.2 Structures of rank 3 orbits under the actions of the groups generated by two conjugate proper rotational symmetries

Let us investigate structures of orbits under the groups generated by two conjugate rotational symmetries. We assume that $R(\mathcal{M})$ contains a proper rotational symmetry of order $p > 2$.

Proposition 5.2 Let $\mathcal{M}$ be a simple rank 3 oriented matroid and G the cyclic group generated by a proper rotational symmetry $\sigma \in R(\mathcal{M})$ of order $p (> 2)$. For distinct rank 3 orbits $X := \{x, \sigma(x), \dots, \sigma^{p-1}(x)\}$, $Y := \{y, \sigma(y), \dots, \sigma^{p-1}(y)\}$ under the action of G, if $\tau \cdot X = Y$ for $\tau \in G(\mathcal{M})$ then we have $\sigma\tau = \tau\sigma$ or $\tau^2 = e, (\tau\sigma)^2 = e$. In the first case, the oriented matroid $\mathcal{M}|_{X \cup Y}$ is a relabeling of the alternating matroid $A_{3,2|X|}$. In the latter case, the reorientation ${}_{-Y}\mathcal{M}|_{X \cup Y}$ is a relabeling of the alternating matroid $A_{3,2|X|}$.

Proof: If $X = Y$, the proposition is trivial. We prove the proposition for the case $X \neq Y$ by contradiction. First of all, note that $(\tau\sigma\tau^{-1})|_Y$ is a rotational symmetry of $\mathcal{M}|_Y$. Therefore, there exists $l \in \mathbb{N}$ such that $(\tau\sigma\tau^{-1})|_Y = \sigma^l|_Y$ and thus $\tau\sigma\tau^{-1} = \sigma^l$. First, we assume that $l = 1$ or -1, and prove that $\mathcal{M}|_{X \cup Y}$ or ${}_{-Y}\mathcal{M}|_{X \cup Y}$ is a relabeling of the alternating matroid $A_{3,2|X|}$.

Without of loss of generality, we suppose that $x, \sigma(x), \sigma^2(x), \dots, \sigma^{p-1}(x)$ define an alternating matroid in this order i.e., $\chi(\sigma^i(x), \sigma^j(x), \sigma^k(x)) = +$ for all $0 \le i < j < k \le p-1$, where χ is a chirotope of $\mathcal{M}|_X$. In this setting, $\{x, \sigma(x)\}$ determines a facet of $\mathcal{M}|_X$. Also, note that $\{\tau(x), \sigma(\tau(x))\}$ $(= \tau \cdot \{x, \sigma^{-1}(x)\}$ or $\tau \cdot \{x, \sigma(x)\})$ defines a facet of $\mathcal{M}|_Y$.

From now on, let us write $\mathcal{N} := \mathcal{M}|_{X \cup Y}$ and $x_1 := x, x_2 := \sigma(x), \dots, x_p := \sigma^{p-1}(x)$. As a first step, we prove that $\mathcal{N}$ or ${}_{-Y}\mathcal{N}$ is acyclic. Let us consider the cocircuit V of $\mathcal{N}$ such that

$$V(x_1) = V(x_2) = 0 \text{ and } V(e) = + \text{ for all } e \in X \setminus \{x_1, x_2\}$$

and write $V_0 := V, V_1 := \sigma(V), \dots, V_{p-1} := \sigma^{p-1}(V)$.



Figure 3: Situations

If $V_0(e) = +$ for all $e \in Y$ or $V_0(e) = -$ for all $e \in Y$, it is clear that $\mathcal{N}$ or ${}_{-Y}\mathcal{N}$ is acyclic. Otherwise, it holds that

$$V_0(e_0) = \cdots = V_{l_1}(e_0) = +, V_{l_1+1}(e_0) = \cdots = V_{l_2}(e_0) = -, V_{l_2+1}(e_0) = \cdots = V_{p-1}(e_0) = +$$

for some $0 \leq l_1 < l_2 \leq p-1$ and $e_0 \in Y$ (Case (A)) or

$$V_0(e_0) = \cdots = V_{l_1}(e_0) = -, V_{l_1+1}(e_0) = \cdots = V_{l_2}(e_0) = +, V_{l_2+1}(e_0) = \cdots = V_{p-1}(e_0) = -$$

for some $0 \leq l_1 < l_2 \leq p-1$ and $e_0 \in Y$ (Case (B)), by the property rank 3 alternating matroids (Proposition 7.1 in Appendix). Here, we are assuming that $V_i(e_0) \neq 0$ for $i = 0, \dots, p-1$ for the sake of simplicity. All the discussion below also applies to the case where $V_i(e_0) = 0$ for some i.

First, let us consider Case (A). Without loss of generality (by relabeling $V_0, \dots, V_{p-1}$ appropriately if necessary), we can assume that

$$V_0(e_0) = \cdots = V_{l_1}(e_0) = +, V_{l_1+1}(e_0) = \cdots = V_{p-1}(e_0) = -.$$

If $l_1 = p-1$, then $\mathcal{N}$ is clearly acyclic. In the following, we assume that $0 \leq l_1 \leq p-2$. Applying σ^i to the above relation, we obtain

$$V_0(e_i) = \cdots = V_{i-1}(e_i) = -, V_i(e_i) = \cdots = V_{i+l_1}(e_i) = +, V_{i+l_1+1}(e_i) = \cdots = V_{p-1}(e_i) = -$$

if $1 \leq i \leq p-1-l_1$ and

$$V_0(e_i) = \cdots = V_{i+l_1-p}(e_i) = +, V_{i+l_1-p+1}(e_i) = \cdots = V_{i-1}(e_i) = -, V_i(e_i) = \cdots = V_{p-1}(e_i) = +$$

if $p-1-l_1 < i < p$. Therefore, it holds that

$$V_0(e_0) = +, V_0(e_1) = \cdots = V_0(e_{p-1-l_1}) = -, V_0(e_{p-l_1}) = \cdots = V_0(e_{p-1}) = +.$$

We also see that

$$\begin{aligned}
&V_{p-1}(e_0) = \cdots = V_{p-1}(e_{p-2-l_1}) = -, V_{p-1}(e_{p-1-l_1}) = \cdots = V_{p-1}(e_{p-1}) = +,\\
&V_{l_1}(e_0) = \cdots = V_{l_1}(e_{l_1}) = +, V_{l_1}(e_{l_1+1}) = \cdots = V_{l_1}(e_{p-1}) = -,\\
&V_{l_1+1}(e_0) = -, V_{l_1+1}(e_1) = \cdots = V_{l_1+1}(e_{l_1+1}) = +, V_{l_1+1}(e_{l_1+2}) = \cdots = V_{l_1+1}(e_{p-1}) = -.
\end{aligned}$$

Applying vector elimination to V_0, V_{p-1} and e_0 (resp. V_{l_1}, V_{l_1+1} and e_0), we obtain a covector W_1 (resp. W_2) such that

$$W_1(e_0) = 0, W_1(e_1) = \cdots = W_1(e_{p-2-l_1}) = -, W_1(e_{p-l_1}) = \cdots = W_1(e_{p-1}) = +, \text{ and}$$

$$W_2(e_0) = 0, W_2(e_1) = \cdots = W_2(e_{l_1}) = +, W_2(e_{l_1+2}) = \cdots = W_2(e_{p-1}) = -.$$

If $1 \leq l_2 \leq p-3$, then apply vector elimination to W_1, W_2 and e_1. Since $\{e_0, e_1\}$ defines a facet of $\mathcal{M}|_Y$, we obtain a covector W_3 such that

$$W_3(e_0) = W_3(e_1) = 0,\ W_3(e) = + \text{ for all } e \in X,\ W_3(e) = + \text{ for all } Y \setminus \{e_0, e_1\}, \text{ or}$$

$$W_3(e_0) = W_3(e_1) = 0,\ W_3(e) = + \text{ for all } e \in X,\ W_3(e) = - \text{ for all } Y \setminus \{e_0, e_1\}.$$

Clearly, it holds that $W_3(e) = +$ for all $e \in X$. By considering $W_3 \circ \sigma(W_3) \circ \cdots \circ \sigma^{p-1}(W_3)$, the oriented matroid $\mathcal{N}$ or ${}_{-Y}\mathcal{N}$ is proved to be acyclic. If $l_1 = p-2$ and $W_2(e_{p-1}) \geq 0$, then $\mathcal{N}$ is acyclic because of the positive covector $W_2 \circ \sigma(W_2) \circ \cdots \circ \sigma^{p-1}(W_2)$. If $l_1 = p-2$ and $W_2(e_{p-1}) = -$, then apply vector elimination to W_1, W_2 and e_{p-1}. Since $\{e_0, e_{p-1}\}$ defines a facet of $\mathcal{M}|_Y$, we obtain a covector W_4 such that

$$W_4(e_0) = W_4(e_{p-1}) = 0,\ W_4(e) = + \text{ for all } e \in X,\ W_4(e) = + \text{ for all } Y \setminus \{e_0, e_{p-1}\}, \text{ or}$$

$$W_4(e_0) = W_4(e_{p-1}) = 0,\ W_4(e) = + \text{ for all } e \in X,\ W_4(e) = - \text{ for all } Y \setminus \{e_0, e_{p-1}\}.$$

Clearly, it holds that $W_4(e) = +$ for all $e \in X$. Therefore, $\mathcal{N}$ or ${}_{-Y}\mathcal{N}$ is acyclic because of the positive covector $W_4 \circ \sigma(W_4) \circ \cdots \circ \sigma^{p-1}(W_4)$.

The above discussion also applies to the case where $V_i(e_0) = 0$ for some i and Case (B). Therefore, it can be concluded that the oriented matroid $\mathcal{N}$ or ${}_{-Y}\mathcal{N}$ is acyclic.

Now, let us investigate structure of $\mathcal{N}$ under the conclusion that $\mathcal{N}$ or ${}_{-Y}\mathcal{N}$ is acyclic. If $\mathcal{N}$ is acyclic, all elements of $\mathcal{N}$ are extreme points by transitivity (in $\mathcal{M}$) and thus $\mathcal{N}$ is a relabeling of the alternating matroid $A_{3,|X|+|Y|}$.

Let us consider the case when ${}_{-Y}\mathcal{N}$ is acyclic. Then, for every $x_a \in X$ there is the covector Z_a such that

$$Z_a(x_a) = 0, Z_a(e) = + \text{ for all } e \in X \setminus \{x_a\} \text{ and } Z_a(f) = - \text{ for all } f \in Y,$$

and for every $y_b \in Y$ there is a covector Z'_b such that

$$Z'_b(y_b) = 0,\ Z'_b(e) = - \text{ for all } e \in X \text{ and } Z'_b(f) = + \text{ for all } f \in Y \setminus \{y_b\}.$$

Thus, the reorientation ${}_{-Y}\mathcal{N}$ is a matroid polytope of rank 3, i.e., a relabeling of the alternating matroid $A_{3,|X|+|Y|}$. Therefore, the inseparability graph of $\mathcal{N}$ is the $(|X|+|Y|)$-cycle. $\tau|_{X\cup Y}$ and $\sigma|_{X\cup Y}$ are also symmetries of the inseparability graph (for inseparability graphs, see [9]). Therefore, $(\tau\sigma)|_{X\cup Y} \neq (\sigma\tau)|_{X\cup Y}$ leads $\tau^2|_{X\cup Y} = e_{X\cup Y}$ and $(\tau\sigma)^2|_{X\cup Y} = e|_{X\cup Y}$. By Corollary 4.3, we have $\tau^2 = e, (\tau\sigma)^2 = e$.

In the above, we saw that $\mathcal{N}$ is a relabeling of the alternating matroid $A_{3,2|X|}$ if $\tau\sigma\tau^{-1} = \sigma$ and that ${}_{-Y}\mathcal{N}$ is a relabeling of the alternating matroid $A_{3,2|X|}$ if $\tau\sigma\tau^{-1} = \sigma^{-1}$. In the remaining part, we assume that $\tau\sigma\tau^{-1} = \sigma^l$ for $2 \leq l \leq p-2$ and lead a contradiction. Note that $\tau \cdot \{x, \sigma(x)\} = \{\tau(x), \sigma^l(\tau(x))\}$ must be a facet of $\mathcal{M}|_Y$ in this case.

Similarly to the above discussion, let us consider the cocircuit V of $\mathcal{N}$ such that

$$V(x_1) = V(x_2) = 0 \text{ and } V(e) = + \text{ for all } e \in X \setminus \{x_1, x_2\}$$



Figure 4: The case where ${}_{-Y}\mathcal{N}$ is acyclic (represented as a signed point configuration. White points: positive points, black points: negative points.)

and write $V_0 := V, V_1 := \sigma(V), \ldots, V_{p-1} := \sigma^{p-1}(V)$. If $V_0(e) = +$ for all $e \in Y$ or $V_0(e) = -$ for all $e \in Y$, then $\mathcal{N}$ or ${}_{-Y}\mathcal{N}$ is acyclic and thus $l = 1$ or -1, a contradiction. Then, assuming that $V_0(e) \neq 0$ for all $e \in Y$ similarly to the above, we have

$$V_0(e_0) = V_1(e_0) = \cdots = V_k(e_0) = -, V_{k+1}(e_0) = \cdots = V_{p-1}(e_0) = +, \text{ or}$$

$$V_0(e_0) = V_1(e_0) = \cdots = V_k(e_0) = +, V_{k+1}(e_0) = \cdots = V_{p-1}(e_0) = -$$

for $e_0 \in Y$, $e_1 := \sigma(e_0), \ldots, e_{p-1} := \sigma^{p-1}(e_0)$ and $0 \leq k \leq p-2$ (by relabeling $V_0, \ldots, V_{p-1}$ appropriately). First, let us consider the first case. By the same argument as the above, we have

$$V_k(e_0) = V_k(e_1) = \cdots = V_k(e_k) = -, V_k(e_{k+1}) = \cdots = V_k(e_{p-1}) = +, \text{ and}$$

$$V_{k+1}(e_0) = +, V_{k+1}(e_1) = V_{k+1}(e_2) = \cdots = V_{k+1}(e_{k+1}) = -, V_{k+1}(e_{k+2}) = \cdots = V_{k+1}(e_{p-1}) = +.$$

Applying vector elimination to V_k, V_{k+1} and e_0, we obtain a covector W such that

$$W(e_0) = 0, W(e_1) = W(e_2) = \cdots = W(e_k) = -, W(e_{k+2}) = W(e_{k+3}) = \cdots = W(e_{p-1}) = +.$$

Since $e_0, \sigma^l(e_0), \sigma^{2l}(e_0), \ldots, \sigma^{(p-1)l}(e_0)$ define an alternating matroid in this order, the following must hold by the property of rank 3 alternating matroids (Proposition 7.2 in Appendix).

$$\begin{cases} \{\sigma^l(e_0), \sigma^{2l}(e_0), \ldots, \sigma^{kl}(e_0)\} = \{e_1, \ldots, e_k\} \text{ or } \{e_{k+1}, \ldots, e_{p-1}\} \\ \text{ (i.e., } \{l, 2l, \ldots, kl\} \bmod p = \{1, \ldots, k\} \text{ or } \{k+1, \ldots, p-1\}) \\ \text{and} \\ \{\sigma^{(k+1)l}(e_0), \sigma^{(k+2)l}(e_0), \ldots, \sigma^{(p-1)l}(e_0)\} = \{e_1, \ldots, e_k\} \text{ or } \{e_{k+1}, \ldots, e_{p-1}\} \\ \text{ (i.e., } \{(k+1)l, (k+2)l, \ldots, (p-1)l\} \bmod p = \{1, \ldots, k\} \text{ or } \{k+1, \ldots, p-1\}) \end{cases}$$

or

$$\begin{cases} \{\sigma^l(e_0), \sigma^{2l}(e_0), \ldots, \sigma^{(k+1)l}(e_0)\} = \{e_1, \ldots, e_{k+1}\} \text{ or } \{e_{k+2}, \ldots, e_{p-1}\} \\ \text{ (i.e., } \{l, 2l, \ldots, (k+1)l\} \bmod p = \{1, \ldots, k+1\} \text{ or } \{k+2, \ldots, p-1\}) \\ \text{and} \\ \{\sigma^{(k+2)l}(e_0), \sigma^{(k+3)l}(e_0), \ldots, \sigma^{(p-1)l}(e_0)\} = \{e_1, \ldots, e_{k+1}\} \text{ or } \{e_{k+2}, \ldots, e_{p-1}\} \\ \text{ (i.e., } \{(k+2)l, (k+3)l, \ldots, (p-1)l\} \bmod p = \{1, \ldots, k+1\} \text{ or } \{k+2, \ldots, p-1\}) \end{cases}$$

It is impossible since $2 \leq l \leq p-2$. A contradiction also occurs in the latter case. $\square$

Proposition 5.3 Let $\mathcal{M}$ be a simple oriented matroid of rank 3 on a ground set E, σ a proper rotational symmetry of $\mathcal{M}$ of order $p(>2)$ and $\tau \in G(\mathcal{M})$ a rotational symmetry of order q. If $\sigma\tau = \tau\sigma$, then σ and τ are in the same cyclic group.

PROOF: Let $X := \{x, \sigma(x), \ldots, \sigma^{p-1}(x)\}$ be a rank 3 orbit and $X_2 := X \cup \tau(X), \ldots, X_q := X \cup \tau(X) \cup \cdots \cup \tau^{q-1}(X)$. Note that $\mathcal{M}|_X$ and $\mathcal{M}|_{X_2}$ are relabelings of alternating matroids. In the following, we prove that $\mathcal{M}|_{X_{i+1}}$ is a reorientation of an alternating matroid under the assumption that $\mathcal{M}|_{X_i}$ is a relabeling of an alternating matroid.

Without loss of generality, we assume that $\{x, \sigma(x)\}$ (resp. $\{x, \tau(x)\}$) defines a facet of $\mathcal{M}|_X$ (resp. $\mathcal{M}|_{X_i}$). It is sufficient to show that $\mathcal{M}|_{X_{i+1}}$ or ${}_{-\tau^i(X)}\mathcal{M}|_{X_{i+1}}$ is acyclic. Let V be the cocircuit of $\mathcal{M}|_{X_i}$ such that $V(x) = V(\tau(x)) = 0$ and $V(e) = +$ for all $e \in X \cup \tau(X) \setminus \{x, \tau(x)\}$.

(I) $V(e) = +$ for all $e \in \tau^i(X)$.

Since $V \circ \sigma(V) \circ \cdots \circ \sigma^{p-1}(V)$ is the positive covector, the oriented matroid $\mathcal{M}|_{X_{i+1}}$ is acyclic.

(II) $V(e_0) = -$ for some $e_0 \in \tau^i(X)$.

Let us label the elements of X_i by $x_0 (:= x), x_1 \ldots, x_{pi-1}$ so that they define an alternating matroid in this order and set $e_1 := \sigma(e_0), \ldots, e_{p-1} := \sigma^{p-1}(e_0)$. Note that $x_i = \sigma(x), x_{2i} = \sigma^2(x), \ldots, x_{(p-1)i} = \sigma^{p-1}(x)$. Let V_k be the cocircuits of $\mathcal{M}|_{X_{i+1}}$ such that

$$V_k(x_k) = V_k(x_{k+1}) = 0, V_k(e) = + \text{ for all } e \in Y_i \setminus \{x_k, x_{k+1}\},$$

for $k = 1, \ldots, pi-1$, where $x_{pi} := x_0$. Then there exist $l_1, l_2 (l_2 < l_1) \in \mathbb{N}$ such that

$$\begin{cases} V_{l_1}(e_0) = V_{l_1+1}(e_0) = V_{pi-1}(e_0) = V_0(e_0) = V_1(e_0) = \cdots = V_{l_2-1}(e_0) = +, \\ V_{l_2}(e_0) = V_{l_2+1}(e_0) = \cdots = V_{l_1-1}(e_0) = - \end{cases} \tag{1}$$

or

$$\begin{cases} V_{l_1}(e_0) = V_{l_1+1}(e_0) = V_{pi-1}(e_0) = V_0(e_0) = V_1(e_0) = \cdots = V_{l_2-1}(e_0) = -, \\ V_{l_2}(e_0) = V_{l_2+1}(e_0) = \cdots = V_{l_1-1}(e_0) = + \end{cases} \tag{2}$$

by Proposition 7.1 in Appendix. Let us first consider Case (1). Without loss of generality, we can assume that $0 \leq l_2 \leq i-1$. For $k = 1, \ldots, p-1$, let us apply σ^k. Then, we obtain

$$\begin{cases} V_{l_1+ki}(e_k) = V_{l_1+ki+1}(e_k) = V_{pi-1}(e_k) = V_0(e_k) = V_1(e_k) = \cdots = V_{l_2+ki-1}(e_k) = +, \\ V_{l_2+ki}(e_k) = V_{l_2+ki+1}(e_k) = \cdots = V_{l_1+ki-1}(e_k) = - \end{cases}$$

if $ki + l_1 < pi$. On the other hand, we have

$$\begin{cases} V_{l_1+ki-pi}(e_k) = V_{l_1+ki-pi+1}(e_k) = \cdots = V_{l_2+ki-1}(e_k) = +, \\ V_0(e_k) = V_1(e_k) = \cdots = V_{l_1+ki-pi-1}(e_k) = -, V_{l_2+ki}(e_k) = V_{l_2+ki+1}(e_k) = \cdots = V_{pi-1}(e_k) = - \end{cases}$$

if $ki + l_1 \geq pi$. Therefore, the following holds.

$$\begin{cases} V_{l_2-1}(e_0) = \cdots = V_{l_2-1}(e_{p-\lfloor \frac{l_1}{i} \rfloor}) = +, \ V_{l_2-1}(e_{p-\lfloor \frac{l_1}{i} \rfloor+1}) = \cdots = V_{l_2-1}(e_{p-1}) = -, \\ V_{l_2}(e_0) = -, \ V_{l_2}(e_1) = \cdots = V_{l_2}(e_{p-\lfloor \frac{l_1}{i} \rfloor+1}) = +, \ V_{l_2}(e_{p-\lfloor \frac{l_1}{i} \rfloor+2}) = \cdots = V_{l_2}(e_{p-1}) = -, \end{cases}$$

$$\begin{cases} V_{l_3}(e_0) = \cdots = V_{l_3}(e_{\lfloor \frac{l_1}{i} \rfloor-2}) = -, \ V_{l_3}(e_{\lfloor \frac{l_1}{i} \rfloor-1}) = \cdots = V_{l_3}(e_{p-1}) = +, \\ V_{l_3+1}(e_0) = +, \ V_{l_3+1}(e_1) = \cdots = V_{l_3+1}(e_{\lfloor \frac{l_1}{i} \rfloor-1}) = -, \ V_{l_3+1}(e_{\lfloor \frac{l_1}{i} \rfloor}) = \cdots = V_{l_3+1}(e_{p-1}) = +, \end{cases}$$

where $l_3 := l_2 - 1 + (\lfloor \frac{l_1}{i} \rfloor - 1)i$.



Figure 5: Situations

(a) $l_1 \leq i$.
The oriented matroid $\mathcal{M}|_{X_{i+1}}$ is acyclic because of the positive vector $V_{l_2} \circ \sigma(V_{l_2}) \circ \cdots \circ \sigma^{p-1}(V_{l_2})$.

(b) $l_1 > i$.
Let us apply vector elimination to V_{l_2-1}, V_{l_2} and e_0 (resp. V_{l_3}, V_{l_3+1} and e_0), and obtain a covector W_1 (resp. W_2) such that

$$W_1(e_0) = W_1(x_{l_2}) = 0, W_1(e) = + \text{ for all } e \in X_i \setminus \{x_{l_2}\}.$$

$$W_2(e_0) = W_2(x_{l_3+1}) = 0, W_2(e) = + \text{ for all } e \in X_i \setminus \{x_{l_3+1}\}.$$

By the same argument as the proof of Proposition 5.2, the oriented matroid $\mathcal{M}|_{X_{i+1}}$ or ${}_{-\tau^i(X)}\mathcal{M}|_{X_{i+1}}$ is proved to be acyclic. The same argument applies to Case (2). Therefore, the oriented matroid $\mathcal{M}|_{X_{i+1}}$ or ${}_{-\tau^i(X)}\mathcal{M}|_{X_{i+1}}$ is a relabeling of the alternating matroid $A_{3,|X_{i+1}|}$. Since σ and τ are also symmetries of ${}_{-\tau^i(X)}\mathcal{M}$, we can continue this argument until $\mathcal{M}|_{X_q}$ is proved to be a reorientation of the alternating matroid $A_{3,|X_q|}$. Therefore, the inseparability graph of $\mathcal{M}|_{X_q}$ is the $|X_q|$-cycle and this leads that σ and τ are in the same cyclic group. $\square$

6 Rotational symmetry groups of simple acyclic oriented matroids of rank 4

In this section, we classify rotational symmetry groups of simple acyclic oriented matroids of rank 4 using the fixed point theorems and the results in the previous section.

6.1 Condition for a subgroup of $R(\mathcal{M})$ to be a cyclic group

First of all, let us investigate, as the simplest case, a condition that a subgroup of $R(\mathcal{M})$ is a cyclic group.

Proposition 6.1 Let $\mathcal{M} = (E, \{\chi, -\chi\})$ be a simple acyclic oriented matroid of rank 4 and $\sigma \in R(\mathcal{M})$. A subgroup $G \subseteq R(\mathcal{M})$ is a cyclic group if and only if $rank(\mathcal{M}|_{G \cdot x}) \leq 3$ for all $x \in E$.

Proof: ($\Leftarrow$) If $rank(\mathcal{M}|_{G\cdot x}) = 1$ for all $x \in E$, then $G = \{e\}$ and thus G is a cyclic group. If $rank(\mathcal{M}|_{G\cdot x}) \leq 2$ for all $x \in E$ and $G \neq \{e\}$, then $|G| = 2$. This means that G is a cyclic group. If there exists $x \in E$ such that $rank(\mathcal{M}|_{G\cdot x}) = 3$, the group $G' := \{\tau|_{G\cdot x} \mid \tau \in G\}$ is a subgroup of rank 3 simple acyclic oriented matroid $\mathcal{M}|_{G\cdot x}$ and thus a cyclic group. By Corollary 4.3, G is also a cyclic group.
($\Rightarrow$) Let us consider the oriented matroid $\mathcal{N} := \mathcal{M}|_{G\cdot x}$. Then all elements of $\mathcal{N}$ are extreme points by transitivity. We prove a contradiction by assuming $rank(\mathcal{N}) = 4$.

Let us consider a single element extension $\mathcal{N} \cup q$ of $\mathcal{N}$ with a fixed point q using Theorem 4.5. Then contract $\mathcal{N} \cup q$ by q and obtain $\mathcal{N}' := (\mathcal{N} \cup q)/q$. Note that $\sigma \in R(\mathcal{N}')$. By the choice of q, the oriented matroid $\mathcal{N}'$ is not acyclic (and loopless). Therefore, $\mathcal{N}'$ is not simple by Proposition 5.1. Note that no two distinct elements are parallel and no three distinct elements are antiparallel in $\mathcal{N}'$ by the convexity of $\mathcal{N}$. From the above information and Proposition 5.1, each element of $\mathcal{N}'$ has exactly one distinct antiparallel element and no parallel element as in Figure 6.



Figure 6: non-simple case (without loops)

Let $a : E \to E$ be the map that takes each element of $\mathcal{N}'$ to its antiparallel element. Let us take $x, y, z \in E$ such that $\chi(x, y, z) \neq 0$ Note that $\sigma(a(x)) = a(\sigma(x))$ (since $\sigma(x)$ and $\sigma(a(x))$ must be antiparallel). Let $k > 0$ be the smallest number such that $\sigma^k(x) = a(x)$. Because of the transitivity, this number is common for all elements in E. Thus

$$\sigma^k \cdot \chi(x, y, z) = \chi(a(x), a(y), a(z)) = -\chi(x, y, z).$$

It contradicts to the fact that σ^k is a rotational symmetry of $\mathcal{N}'$. This implies that the case $rank(\mathcal{N}) = 4$ cannot occur. As a conclusion, it holds that $rank(\mathcal{M}|_{G\cdot x}) \leq 3$ for all $x \in E$. $\square$

6.2 Preparation for classification of rotational symmetry groups

In group theory, the standard approach to classify finite subgroups of the special orthogonal group $SO(3)$ is to consider point sets where each non-trivial symmetry fixes exactly two points (see [14]). However, it seems hard to follow the approach in the setting of oriented matroids directly. To follow the excellent classical approach in group theory, our first goal is to construct a certain set on which $R(\mathcal{M})$ acts and the number of fixed elements under each element of $R(\mathcal{M})$ is *uniform*. We will construct a desired set, based on orbits under maximal cyclic subgroups of $R(\mathcal{M})$. First, let us note that each element of $R(\mathcal{M}) \setminus \{e\}$ belongs to exactly one maximal cyclic subgroup of $R(\mathcal{M})$.

Proposition 6.2 Let $\mathcal{M}$ be a simple acyclic oriented matroid of rank 4 on a ground set E and G_1, G_2 distinct maximal cyclic subgroups of $R(\mathcal{M})$. Then $G_1 \cap G_2 = \{e\}$.

Proof: Let σ (resp. τ) be a generator of G_1 (resp. G_2), $H_1 := G_1 \cap G_2$ and H_2 the group generated by $G_1 \cup G_2$.

If $|H_1| \geq 3$, there exists $x \in E$ such that $rank(H_1 \cdot x) = 3$. Thus it holds that $Flat(G_1 \cdot x) = Flat(G_2 \cdot x)$. Therefore, the restriction $\mathcal{M}|_{H_2 \cdot x}$ is a rank 3 matroid polytope. This leads that there exists a cyclic group $G' \supseteq G_1, G_2$, a contradiction.

If $|H_1| = 2$, there exists $y \in E$ such that $rank(H_1 \cdot y) = 2$. Since $|G_1|, |G_2| \geq 3$, it holds that $rank(G_1 \cdot y) = rank(G_2 \cdot y) = 3$. Let $2a, 2b$ be the orders of σ and τ respectively. Then, we have $\sigma^a = \tau^b$. Note that $\sigma^a \tau = \tau \sigma^a$. In the flat $Flat(G_2 \cdot y)$, the elements $\tau(y)$ and $\tau^b(\tau(y))$ are on the other sides of $Flat(\{y, \tau^b(y)\})$ $(= Flat(\{y, \sigma^a(y)\}))$. Therefore, the elements $\tau(y)$ and $\tau^b(\tau(y)) = \sigma^a(\tau(y))$ are on the other sides of $Flat(G_1 \cdot y)$ in $\mathcal{M}$. This implies that σ^a is a reflection symmetry of $\mathcal{M}/X$. Since σ^a is a rotational symmetry of $\mathcal{M}$ and $\mathcal{M}|_X$ ($\mathcal{M}|_X$ is acyclic and thus σ must be a rotational symmetry of $\mathcal{M}|_X$ because its order is greater than 2), it is a contradiction. $\square$

Therefore, if we can construct a set such that a certain group action of $R(\mathcal{M})$ on it is defined and the stabilizer subgroup of each element in $R(\mathcal{M})$ is a maximal cyclic subgroup of $R(\mathcal{M})$, it has a desired property. In the following, let us write G_σ to denote the maximal cyclic subgroup of $R(\mathcal{M})$ that contains $\sigma \in R(\mathcal{M})$. G_σ can be can be classified into two types i.e., $G_\sigma \in G^I \cup G^{II}$, where G^I and G^{II} are defined as follows.

$$\begin{aligned} G^I &:= \{G_\sigma \mid \sigma \in R(\mathcal{M}),\ rank(G_\sigma \cdot x) = 1 \text{ or } 3 \text{ for all } x \in E\}, \\ G^{II} &:= \{G_\sigma \mid \sigma \in R(\mathcal{M}),\ rank(G_\sigma \cdot x) = 1 \text{ or } 2 \text{ for all } x \in E\}. \end{aligned}$$

Note that $G_\sigma \in G^{II}$ if and only if $|G_\sigma| = 2$.

(I) Constructing a suitable set for G^I

The first idea may be to consider the set $S(\sigma) := \{G_\sigma \cdot x \mid x \in E\}$ for $\sigma \in R(\mathcal{M})$. However, there is possibility that $S(\sigma)$ is fixed by an element $\tau \in R(\mathcal{M}) \setminus G_\sigma$. The following proposition gives a necessary condition that such an event occurs (in a more general setting).

Proposition 6.3 Let $\mathcal{M}$ be a simple acyclic oriented matroid of rank 4, G a cyclic subgroup of $R(\mathcal{M})$ generated by $\sigma (\neq e)$ and X_1, X_2 orbits under G with $rank(X_1) = 3, rank(X_2) = 3$. For $\tau \in R(\mathcal{M}) \setminus G$, if $\tau \cdot X_1 = X_2$ and $\sigma\tau \neq \tau\sigma$, then $\tau^2 = e$ and $(\sigma\tau)^2 = e$. If $\tau \cdot X_1 = X_2$ and $\sigma\tau = \tau\sigma$, then σ and τ are in the same cyclic subgroup.

Proof: Let H be the group generated by σ and τ, and $X := H \cdot x$ for $x \in X_1$. Note that $X \supseteq X_1, X_2$.

(i) $\sigma\tau \neq \tau\sigma$.

(i-a) $rank(X) = 3$.

First of all, note that every rotational symmetry σ induces a (not necessarily rotational) symmetry of $\mathcal{M}|_X$. Since $\mathcal{M}$ is acyclic, the restriction $\mathcal{M}|_X$ is also acyclic. By transitivity, the oriented matroid $\mathcal{M}|_X$ is a rank 3 matroid polytope, i.e., a relabeling of the rank 3 alternating matroid $A_{3,|X|}$. If $(\tau\sigma)|_X = (\tau\sigma)|_X$, then $\tau|_X$ and $\sigma|_X$ are in the same cyclic subgroup. This implies that σ and τ are in the same cyclic subgroup by Corollary 4.3. Therefore, we have $(\tau\sigma)|_X \neq (\tau\sigma)|_X$. This implies that $\tau|_X$ is a reflection symmetry of $\mathcal{M}|_X$ and thus $\tau^2|_X = e$ and $(\sigma\tau)^2|_X = e$. Therefore, we have $\tau^2 = e, (\sigma\tau)^2 = e$ by Corollary 4.3.

(i-b) $rank(X) = 4$.

First of all, note that $(\sigma\tau)|_X \neq (\tau\sigma)|_X$ because $(\sigma\tau)|_X = (\tau\sigma)|_X$ implies $\sigma\tau = \tau\sigma$. By transitivity, $\mathcal{M}|_X$ is a rank 4 matroid polytope. Let us consider $\mathcal{N} := (\mathcal{M}|_X \cup q)/q$, where q is the fixed point obtained by Theorem 4.5. Every rotational symmetry σ induces a (not necessarily rotational) symmetry of $\mathcal{N}$.

If $\mathcal{N}$ is simple, then $(\tau\sigma)^2 = e$, $\tau^2 = e$ by Proposition 5.2. Thus we assume that $\mathcal{N}$ is non-simple. By a similar argument to the proof of Proposition 6.1, each element of $\mathcal{N}'$ has exactly one distinct antiparallel element and no distinct parallel elements as in Figure 6.

Let $a : X \to X$ be the map that takes each element of X to its antiparallel element. The permutation a is a reflection symmetry of $\mathcal{N}$ (and $\mathcal{M}$). Note that $a(\sigma(x)) = \sigma(a(x))$ and $a(\tau(x)) = \tau(a(x))$ for all $x \in X$ and thus $\tau(a(X_1)) = a(X_2)$.

If there exists $x_0 \in X_1$ such that $\tau(x_0) \notin a(X_1)$, then $a(X_1) \cap X_2 = \emptyset$. Thus a rank 3 oriented matroid $\mathcal{N}|_{X_1 \cup X_2}$ is simple. In this case, we have $\tau^2|_{X_1 \cup X_2} = e, (\sigma\tau)^2|_{X_1 \cup X_2} = e$ by Proposition 5.2. Therefore, we have $\tau^2 = e, (\sigma\tau)^2 = e$.

If $X_2 = a(X_1)$, then $a^{-1}\tau$ is a symmetry of $\mathcal{N}|_{X_1}$. If $(a^{-1}\tau)|_{X_1}$ is a rotational symmetry of $\mathcal{N}|_{X_1}$, there exists $k \in \mathbb{N}$ such that $(a^{-1}\tau)|_{X_1} = \sigma^k|_{X_1}$ and thus $a^{-1}\tau = \sigma^k$. This implies that τ is a reflection symmetry of $\mathcal{M}$, a contradiction. Therefore, $(a^{-1}\tau)|_{X_1}$ is a reflection symmetry of $\mathcal{N}|_{X_1}$ and thus there exists $l \in \mathbb{N}$ such that $(a^{-1}\tau)^2|_{X_1} = e$, $(a^{-1}\tau\sigma)^2|_{X_1} = e$. This leads $\tau^2 = e$ and $(\tau\sigma)^2 = e$ by Corollary 4.3.

(ii) $\sigma\tau = \tau\sigma$.

(ii-a) $rank(X) = 3$.

Since $\sigma|_X$ and $\tau|_X$ are in the same cyclic group, the symmetries σ and τ are in the same cyclic subgroup of $R(\mathcal{M})$.

(ii-b) $rank(X) = 4$.

By the same argument as (i-b), the symmetries σ and τ are proved be in the same cyclic subgroup of $R(\mathcal{M})$.

□

We can remedy the bad situation by considering orderings of rank 3 orbits instead of the set of them. Let us pick up rank 3 orbits under the action of G_σ and consider the flats $F_1, \ldots, F_m$ of them. We assume that there is no duplication in this list by removing some of the flats if necessary. Then, consider flat orderings $(F_{p(1)}, \ldots, F_{p(m)})$ that satisfies the condition in the following lemma.

Lemma 6.1 Let $\mathcal{M}$ be a simple acyclic oriented matroid of rank 4 on a ground set E and $G \subseteq R(\mathcal{M})$ a cyclic subgroup of $R(\mathcal{M})$ of order $q > 2$. Consider rank 3 orbits $O_1, \ldots, O_m$ under the action of G and set $F_1 := Flat(O_1), \ldots, F_m := Flat(O_m)$. We assume that there is no duplication in this list by removing some of them if necessary. Suppose $m \geq 2$. Then, there exist exactly two permutations p on $\{1, \ldots, m\}$ that satisfy the following condition. For every $i = 1, \ldots, m$, there exists a covector X of $\mathcal{M}$ such that

$$\begin{cases} X(e) = 0 \text{ for all } e \in F_{p(i)}, \\ X(f) = - \text{ for all } f \in F_{p(1)} \cup \cdots \cup F_{p(i-1)}, \\ X(g) = + \text{ for all } f \in F_{p(i+1)} \cup \cdots \cup F_{p(m)}. \end{cases}$$

Proof: Let S, T, U be arbitrary rank 3 orbits under the action of G and X, Y be cocircuits whose zero sets are $Flat(S), Flat(T)$ respectively.

Claim. For a cocircuit X of $\mathcal{M}$ such that $X(e) = 0$ for all $e \in Flat(S)$,

$$X(f) = + \text{ for all } f \in T \text{ or } X(f) = - \text{ for all } f \in T.$$

Proof of the claim: First, let us assume that there exists $e_0 \in T$ such that $X(e_0) = 0$. This means $e_0 \in Flat(S)$. By transitivity, $X(e) = 0$ for all $e \in T$, a contradiction. Then, assume that there exist $e_1, e_2 \in T$ such that $X(e_1) = +, X(e_2) = -$. Let $\sigma \in G$ be a rotational symmetry such that $\sigma(e_1) = e_2$. By applying vector elimination to $X, \sigma(X)$ and e_2, we obtain a covector Y such that $Y(e) = 0$ for all $e \in T \cup \{e_2\}$. Note that if $Y = 0$, then $\sigma(X) = -X$ is required. If the requirement is satisfied, the symmetry $\sigma|_{E\setminus X}$ is a reflection symmetry of $\mathcal{M}/X$. Since σ and $\sigma|_X$ are rotational symmetries of $\mathcal{M}$ and $\mathcal{M}|_X$ respectively, it is impossible. Therefore, we have $Y \neq 0$, a contradiction. $\square$

Note that the same holds even if S and T are replaced by other orbits on distinct flats.

Now suppose that

$$X(e) = + \text{ for all } e \in \mathit{Flat}(T)$$

and

$$\begin{cases} Y(e) = - \text{ for all } e \in \mathit{Flat}(S), \\ Y(e) = + \text{ for all } e \in \mathit{Flat}(U). \end{cases}$$

To construct a required permutation, we must be able to construct a covector Z such that

$$\begin{cases} Z(e) = + \text{ for all } e \in \mathit{Flat}(S), \\ Z(e) = + \text{ for all } e \in \mathit{Flat}(T), \\ Z(e) = 0 \text{ for all } e \in \mathit{Flat}(U). \end{cases}$$

If $X(e) = -$ for $e \in U$, it is proved to be impossible as follows.

Under the assumption, we first prove that there is a covector W^* such that

$$\begin{cases} W^*(e) = - \text{ for all } e \in \mathit{Flat}(S), \\ W^*(e) = + \text{ for all } e \in \mathit{Flat}(T), \\ W^*(e) = 0 \text{ for all } e \in \mathit{Flat}(U). \end{cases}$$

Apply conformal elimination to X, Y and U and obtain a covector W_1 such that

$$\begin{cases} W_1(e) = - \text{ for all } e \in \mathit{Flat}(S), \\ W_1(e) = + \text{ for all } e \in \mathit{Flat}(T), \\ W_1(u_0) = 0 \text{ for some } u_0 \in U \text{ and } W_1(e) \in \{0, -\} \text{ for all } U \setminus \{u_0\}. \end{cases}$$

Similarly, we obtain a covector W_2 such that

$$\begin{cases} W_2(e) = - \text{ for all } e \in \mathit{Flat}(S), \\ W_2(e) = + \text{ for all } e \in \mathit{Flat}(T), \\ W_2(u_1) = 0 \text{ for some } u_1 \in U \text{ and } W_2(e) \in \{0, +\} \text{ for all } U \setminus \{u_1\}. \end{cases}$$

If $W_1(e) = 0$ for all $e \in U$ or $W_2(e) = 0$ for all $e \in U$, we obtain the required covector W^*.

Otherwise, let us take $\sigma \in G$ such that $\sigma(u_1) = u_0$ and consider $W_3 := \sigma(W_2)$. Then apply conformal elimination to W_1, W_3 and U to obtain a new covector W_4.

$$\begin{cases} W_4(e) = - \text{ for all } e \in \mathit{Flat}(S), \\ W_4(e) = + \text{ for all } e \in \mathit{Flat}(T), \\ W_4(u_0) = W(u_2) = 0 \text{ for some } u_2 \in U \text{ and } W(e) \in \{0, -\} \text{ for all } U \setminus \{u_0, u_2\}. \end{cases}$$

If $W_4(e) = 0$ for all $e \in U$, we obtain the required covector W^*. Otherwise, take $u_3 \in U$ such that $\{u_0, u_3\}$ defines a facet of $\mathcal{M}|_U$ and $W_4(u_3) \neq 0$ (note that one of two elements adjacent to u_0 satisfies this condition. Otherwise, $W_4(e) = 0$ for all $e \in U$). Apply vector elimination to W_3, W_4 and u_3. Then, the obtained covector W_5 satisfies

$$\begin{cases} W_5(e) = - \text{ for all } e \in \mathit{Flat}(S), \\ W_5(e) = + \text{ for all } e \in \mathit{Flat}(T), \\ W_5(u_0) = W_5(u_3) = 0 \text{ and } W_5(e) = + \text{ for all } e \in U \setminus \{u_0, u_3\}. \end{cases}$$

Similarly, we can construct a covector W_6 such that

$$\begin{cases} W_6(e) = - \text{ for all } e \in \mathit{Flat}(S), \\ W_6(e) = + \text{ for all } e \in \mathit{Flat}(T), \\ W_6(u_0) = W_6(u_3) = 0 \text{ and } W_6(e) = - \text{ for all } e \in U \setminus \{u_0, u_3\}. \end{cases}$$

Applying vector elimination to W_5, W_6 and $u_4 \in U \setminus \{u_0, u_3\}$, we obtain the required covector W^*. Here, note that $W^*(e) = 0$ for all $e \in U$ if $W^*(e_1) = W^*(e_2) = W^*(e_3) = 0$ for distinct elements $e_1, e_2, e_3 \in U$.

Then, apply vector elimination to W^*, Z and $s_0 \in S$, and obtain a covector W^{**} such that

$$W^{**}(e) = 0 \text{ for all } e \in \mathit{Flat}(U) \cup \{s_0\} \text{ and } W^{**}(e) = + \text{ for all } e \in \mathit{Flat}(T).$$

This contradicts to the assumption that $\mathit{rank}(U) = 3$.

On the other hand, the required covector Z can be constructed in the same way as the above discussion if $X(e) = +$ for $e \in U$. This completes the proof. □

Let us write these two orderings as $T_1(\sigma)$ and $T_2(\sigma)$ (it is arbitrarily decided which one goes to $T_1(\sigma)$ and $T_2(\sigma)$). If there is only one flat of rank 3 orbit, we denote it by F. Let X_F be a cocircuit with $X_F(F) = 0$. Then, we set $T_1(\sigma) := (X_F^-, F, X_F^+)$ and $T_2(\sigma) := (X_F^+, F, X_F^-)$, where $X_F^+ := \{e \in E \mid X_F(e) = +\}$ and $X_F^- := \{e \in E \mid X_F(e) = -\}$. Then, consider the collection of $T_1(\sigma)$ and $T_2(\sigma)$:

$$S^I := \{T_1(\sigma) \mid \ G_\sigma \in G^I\} \cup \{T_2(\sigma) \mid \ G_\sigma \in G^I\}$$

Let us consider the standard group action of $R(\mathcal{M})$ on S^I. Clearly, the group action is well-defined i.e., $T_i(\sigma_1) = T_i(\sigma_2) \Rightarrow \tau \cdot T_i(\sigma_1) = \tau \cdot T_i(\sigma_2)$ for all $\sigma_1, \sigma_2 \in R(\mathcal{M})$, for $i = 1, 2$. Furthermore, the group action is closed in S^I. Rank 3 orbits are transformed into rank 3 orbits by the following computation:

$$\tau \cdot \{x, \sigma(x), \sigma^2(x), \dots\} = \{\tau(x), \tau\sigma\tau^{-1}(\tau(x)), (\tau\sigma\tau^{-1})^2(\tau(x)), \dots\}$$

for $\sigma, \tau \in R(\mathcal{M})$. Therefore, it holds that $\tau \cdot T_1(\sigma) = T_1(\tau\sigma\tau^{-1})$ and $\tau \cdot T_2(\sigma) = T_2(\tau\sigma\tau^{-1})$, or $\tau \cdot T_1(\sigma) = T_2(\tau\sigma\tau^{-1})$ and $\tau \cdot T_2(\sigma) = T_1(\tau\sigma\tau^{-1})$.

The following lemma assures that there is a one-to-one correspondence between S^I and G^I.

Proposition 6.4 Let σ, τ are generators of $G_\sigma, G_\tau \in G^I$. Then $T_i(\sigma) = T_i(\tau) \Leftrightarrow G_\sigma = G_\tau$, for $i = 1, 2$.

PROOF: ($\Rightarrow$) Let $X := \{x, \sigma(x), \sigma^2(x), \dots, \sigma^{p-1}(x)\}$ be a rank 3 orbit. Note that other orbits in $S(\sigma)$ do not contain x. Thus there exists a rank 3 orbit $\{x, \tau(x), \tau^2(x), \dots, \tau^{p-1}(x)\} = \{x, \sigma(x), \sigma^2(x), \dots, \sigma^{p-1}(x)\}$. Since X is a rank 3 alternating matroid, this implies that $\sigma|_X = \tau^l|_X$ for some $l \in \mathbb{N}$. Since $\mathit{rank}(X) = 3$, we have $\sigma = \tau^l$. The ($\Leftarrow$) part is trivial. □

By the following proposition, the stabilizer subgroup of $T_i(\sigma)$ in $R(\mathcal{M})$ is G_σ for $i = 1, 2$.

Proposition 6.5 $\tau \cdot T_i(\sigma) = T_i(\sigma) \Leftrightarrow \tau \in G_\sigma$, for $i = 1, 2$.

PROOF: Without loss of generality, we assume that σ is a generator of G_σ. It is clear that $\tau \cdot T_i(\sigma) = T_i(\sigma)$ if $\tau \in G_\sigma$. Now let us assume that $\tau \in R(\mathcal{M}) \setminus G_\sigma$ and prove that $\tau \cdot T_i(\sigma) \neq T_i(\sigma)$. Let X be a rank 3 orbit under the action of G_σ. If $\tau \cdot T_i(\sigma) = T_i(\sigma)$, we have $\tau \cdot X = X$. All rotational symmetries of $\mathcal{M}|_X$ are generated by $\sigma|_X$ and thus $\tau|_X$ is not a rotational symmetry of $\mathcal{M}|_X$ but a reflection symmetry. Since $\tau|_X$ is a reflection symmetry of $\mathcal{M}|_X$ and τ is a rotational symmetry of $\mathcal{M}$, $\tau|_{E \setminus X}$ must be a reflection symmetry of $\mathcal{M}/X$. Therefore, the elements x and $\tau(x)$ are on the opposite side of $\mathit{Flat}(X)$ for $x \notin \mathit{Flat}(X)$. This leads $\tau \cdot F \neq F$ for $F \in T_i(\sigma) \setminus \{\mathit{Flat}(X)\}$. Therefore, we have $\tau \cdot T_i(\sigma) \neq T_i(\sigma)$. □

(II) Constructing a suitable set for G^{II}

As a first step, let us consider a set $S_{pre}(\sigma) := \{O \mid O \subseteq E \text{ is an orbit under the action of } G_\sigma\}$ for each $\sigma \in R(\mathcal{M})$ such that $G_\sigma \in G^{II}$ and collect $S_{pre}(\sigma)$ for $\sigma \in G$ such that $G_\sigma \in G^{II}$:

$$S^{II}_{pre} := \{S_{pre}(\sigma) \mid \; G_\sigma \in G^{II}\}$$

Let us consider the standard group action of $R(\mathcal{M})$ on S^{II}_{pre}. Clearly, the group action is well-defined i.e., $S_{pre}(\sigma_1) = S_{pre}(\sigma_2) \Rightarrow \tau \cdot S_{pre}(\sigma_1) = \tau \cdot S_{pre}(\sigma_2)$ for all $\sigma_1, \sigma_2 \in R(\mathcal{M})$. Furthermore, the group action is closed in S by the following computation:

$$\tau \cdot \{x, \sigma(x)\} = \{\tau(x), \tau\sigma\tau^{-1}(\tau(x))\}$$

for $\sigma, \tau \in R(\mathcal{M})$ (note that $\sigma^2 = (\tau\sigma\tau^{-1})^2 = e$).

The following proposition shows that for a rotational symmetry $\sigma \in R(\mathcal{M})$ such that $G_\sigma \in G^{II}$, the stabilizer subgroup of $S_{pre}(\sigma)$ in $R(\mathcal{M})$ is *nearly equal* to G_σ.

Proposition 6.6 Let $\mathcal{M}$ be a simple acyclic oriented matroid of rank 4 on a ground set E and X, Y rank 2 orbits under $\sigma \in R(\mathcal{M})$. If $\tau \cdot S_{pre}(\sigma) = S_{pre}(\sigma)$, then the group H generated by σ and τ is isomorphic to C_2 or $C_2 \times C_2$.

PROOF: Let $X := \{x, \sigma(x)\}$, $Y := \{y, \sigma(y)\}$ and $\tau \cdot X = Y$. If $\tau(x) = y$, $\tau\sigma(x) = \sigma\tau(x)$. If $\tau\sigma(x) = y$, $\sigma\tau\sigma(x) = \tau(x)$. Thus $(\tau\sigma)|_{X\cup Y} = (\sigma\tau)|_{X\cup Y}$ in any cases.

Now assume that $rank(X \cup Y) = 2$. Then, we have $X = Y$ and $\sigma|_X = \tau|_X$. Here, note that

$$rank(Fix(\sigma)) \leq 2, \; rank(Fix(\sigma) \cup X) \leq 3$$

since $\chi(e, f, x, \sigma(x)) = \sigma \cdot \chi(e, f, x, \sigma(x)) = -\chi(e, f, x, \sigma(x))$ for a chirotope χ of $\mathcal{M}$ and $e, f \in Fix(\sigma)$. Thus, there must exist a rank 2 orbit X' (under σ) with $rank(X \cup X') \geq 3$. Let $Y' := \tau \cdot X'$. If $X' = Y'$, then $\tau|_{X\cup X'} = \sigma|_{X\cup X'}$. Since $rank(X \cup X') \geq 3$, we have $\tau = \sigma$ by Corollary 4.3, a contradiction. This implies $rank(X' \cup Y') \geq 3$. Therefore, we can assume that $rank(X \cup Y) \geq 3$ by replacing X, Y by X', Y'.

In this setting, we have $rank(X \cup Y) \geq 3$ and this leads $\sigma\tau = \tau\sigma$ by Corollary 4.3. Let p be the order of τ. If $rank(X \cup Y) = 3$ and $p > 2$, we have $H \simeq C_{2p}$, which contradicts to the maximality assumption of G_σ. If $p = 2$, then $H \simeq C_4$ or $D_4 (\simeq C_2 \times C_2)$ or C_2. If $rank(\mathcal{M}|_{X\cup Y}) = 4$, let us consider the contraction by the fixed point obtained by Theorem 4.5. If $p > 2$, we have $H \simeq C_{2p}$ by the same argument as Proposition 6.3, a contradiction. If $p = 2$, then $H \simeq C_4$ or $C_2 \times C_2$ or C_2. Since $H \simeq C_4$ contradicts to the maximality assumption, we have $H \simeq C_2$ or $C_2 \times C_2$. □

Therefore, the set $S_{pre}(\sigma)$ may be fixed by $\tau \notin G_\sigma$ if σ and τ generate a group isomorphic to $C_2 \times C_2$. The following two propositions show that there cannot be two rotational symmetries with this property.

Proposition 6.7 Let $\mathcal{M}$ be a simple acyclic oriented matroid of rank 4. If there exist $\sigma, \tau, \pi \in R(\mathcal{M})$ of order 2 such that the group generated by σ and τ and that generated by σ and π are isomorphic to $C_2 \times C_2$, then the symmetries σ and $\pi\tau$ are in the same cyclic group, or the group generated by σ, τ and π is isomorphic to $C_2 \times C_2 \times C_2$ or $C_2 \times C_4$.

PROOF: Let p be the order of $\pi\tau$. Note that $\sigma(\pi\tau) = (\pi\tau)\sigma$. If $p > 2$, then σ and τ are in the same cyclic group by Proposition 5.3. If $p = 2$, then the group generated by σ and τ is isomorphic to $C_2 \times C_2$ or C_4. □

Proposition 6.8 Let $\mathcal{M}$ be a simple acyclic oriented matroid of rank 4. Then $R(\mathcal{M})$ does not contain a subgroup G isomorphic to $C_2 \times C_2 \times C_2$ or $C_2 \times C_4$.

Proof: A proof can be done similarly to Proposition 5.3. Let σ, τ, π be generators of G and X be an orbit under the action of G. Then, we have $rank(X) = 3$ or $rank(X) = 4$. If $rank(X) = 4$, consider the contraction by the fixed point. $\square$

Note that the case where G is isomorphic to $C_2 \times C_4$ can be excluded by the maximality assumption.

Based on the above observation, let us define a new set. For a rank 2 orbit $O_0 := \{x, \sigma(x)\}$ such that $O_0 \neq \tau(O_0)$, consider the pairs $(O_0, \tau(O_0))$ and $(\tau(O_0), O_0)$, where τ is a rotational symmetry of order 2 such that the group generated by σ and τ is isomorphic to $C_2 \times C_2$:

$$\begin{aligned} S_1(\sigma) &:= S_{pre}(\sigma) \cup \{(O_0, \tau(O_0))\}, \\ S_2(\sigma) &:= S_{pre}(\sigma) \cup \{(\tau(O_0), O_0)\}. \end{aligned}$$

If there does not exist $\tau \in R(\mathcal{M})$ such that the group generated by σ and τ is isomorphic to $C_2 \times C_2$, we set $S_1(\sigma) = (1, S_{pre}(\sigma))$ and $S_2(\sigma) = (2, S_{pre}(\sigma))$. Then, the stabilizer subgroup of $S_1(\sigma)$ and that of $S_2(\sigma)$ are equal to G_σ. Note that $\tau \cdot S_1(\sigma) = S_2(\sigma)$ and $\tau \cdot S_2(\sigma) = S_1(\sigma)$ if $\tau \in R(\mathcal{M}) \setminus G_\sigma$ satisfies $\tau \cdot S_{pre}(\sigma) = S_{pre}(\sigma)$. Let us define S^{II} as the collection of $S_1(\sigma)$ and $S_2(\sigma)$:

$$S^{II} := \{S_1(\sigma) \mid \sigma \in R(\mathcal{M}), G_\sigma \in G^{II}\} \cup \{S_2(\sigma) \mid \sigma \in R(\mathcal{M}), G_\sigma \in G^{II}\}.$$

The following lemma assures that there is a one-to-one correspondence between S^{II} and G^{II}.

Lemma 6.2 *Let σ, τ be generators of maximal cyclic subgroups of $R(\mathcal{M})$ of order 2. Then $S_i(\sigma) = S_i(\tau) \Rightarrow \sigma = \tau$, for $i = 1, 2$.*

Proof: Let $\{x, \sigma(x)\}$ be a rank 2 orbit under the group generated by σ. Then $\{x, \sigma(x)\} = \{x, \tau(x)\}$. For rank 1 orbit $\{y\}$ $(\sigma(y) = y)$, $\tau(y) = y$. The same applies to all the orbits and thus $\sigma = \tau$. $\square$

6.3 Classification

Now let us classify rotational symmetry groups of simple acyclic oriented matroids of rank 4. From now on, we simply write G instead of $R(\mathcal{M})$.

Consider the group action of G on a set $S := S^I \cup S^{II}$. Let r be the number of orbits in S. First, note the well-known formula (Burnside's lemma):

$$r = \frac{1}{|G|} \sum_{\sigma \in G} f(\sigma),$$

where $f(\sigma)$ is the number of fixed points of σ in S. If σ is the identity, then $f(\sigma) = |S|$ and otherwise $f(\sigma) = 2$.

Lemma 6.3 *$f(\sigma) = 2$ for all $\sigma (\neq e) \in R(\mathcal{M})$.*

Proof: Clearly, it holds that $\sigma \cdot S(\sigma') = S(\sigma')$, where σ' is a generator of a maximal cyclic subgroup of $R(\mathcal{M})$ to which σ belongs if the order of σ' is 2. The same applies to $T_1(\sigma')$ and $T_2(\sigma')$ if the order of σ' is greater than 2. Therefore, we have $f(\sigma) \geq 2$. It is trivial that $f(\sigma) \leq 2$. $\square$

Therefore, we have

$$r = \frac{1}{|G|}\{|S| + 2(|G| - 1)\}.$$

For $x_i \in S_i$, let G_i be the stabilizer subgroup of x_i. Then $|S_i| = \frac{|G|}{|G_i|}$. Note that $|G_i| \geq 2$ and thus

$$r = 2 + \frac{1}{|G|}(|S_1| + \cdots + |S_r| - 2) = 2 - \frac{2}{|G|} + (\frac{1}{|G_1|} + \cdots + \frac{1}{|G_r|}) \leq 2 - \frac{2}{|G|} + \frac{r}{2}.$$

This leads $r < 4$. Also, note that $r \geq 2$. Therefore, $r = 2$ or 3. If $r = 2$, then $|S_1| + |S_2| = 2$ and thus $|S_1| = |S_2| = 1$. Let $s \in S_1$. Then $G \cdot s = s$ and thus G is a cyclic group. If $r = 3$, we have

$$1 < 1 + \frac{2}{|G|} = \frac{1}{|G_1|} + \frac{1}{|G_2|} + \frac{1}{|G_3|}.$$

Therefore, possible types of $(|G_1|, |G_2|, |G_3|, |G|)$ $(|G_1| \geq |G_2| \geq |G_3|)$ are classified into

$$(n, 2, 2, 2n)(n \geq 2), (3, 3, 2, 12), (4, 3, 2, 24), (5, 3, 2, 60).$$

(i) $(|G_1|, |G_2|, |G_3|, |G|) = (n, 2, 2, 2n)$ $(n \geq 2)$.

In this case, we have $|S| = 2n + 2$, $|S_1| = 2$, $|S_2| = |S_3| = n$. Let g, h_1, h_2 be generators of G_1, G_2, G_3 respectively. Then G is generated by g and h_1. Let $\{s, t\} := S_1$.

If $s, t \in S^I$, the flat ordering t must be the reverse ordering of s. Therefore, $h_1 \cdot X_1 = X_2$ for some rank 3 orbits X_1, X_2 for G_1. By Proposition 6.3, we have $h_1^2 = e, (h_1\sigma)^2 = e$. Therefore, G can be regarded as a subgroup of the dihedral group D_{2n}. Since $|G| = 2n$, the group G is, in fact, isomorphic to D_{2n}.

If $s, t \in S^{II}$ and $n \geq 3$, we have $h_1^2 = e, (h_1\sigma)^2 = e$, or h_1 and σ are in the same cyclic subgroup of G. If $s, t \in S^{II}$ and $n = 2$, then $G \simeq C_4$ or $G \simeq D_4$.

(ii) $(|G_1|, |G_2|, |G_3|, |G|) = (3, 3, 2, 12)$.

A group of order 12 is isomorphic to $C_{12}, D_{12}, A_4, C_2 \times C_6$ or Q_{12} (One can check it by using the GAP Small Groups Library [1], for example). Here, Q_{4n} is the bicyclic group defined by $\langle x, y \mid x^2 = y^n, yxy^{-1} = x^{-1}\rangle$. Since G contains two non-conjugate cyclic subgroups of order 3, it holds that $G \not\simeq C_{12}, D_{12}, C_2 \times C_6, Q_{12}$. The remaining possibility is $G \simeq A_4$.

(iii) $(|G_1|, |G_2|, |G_3|, |G|) = (4, 3, 2, 24)$.

A group of order 24 is isomorphic to C_{24}, $C_2 \times C_{12}$, $C_2 \times C_2 \times C_6$, D_{24}, Q_{24}, $C_2 \times D_{12}$, $C_2 \times Q_{12}$, $C_2 \times A_4$, $C_3 \times D_8$, $C_3 \times Q_8$, $C_4 \times D_6$, $SL(2, 3) := \langle a, b, c \mid a^3 = b^3 = c^2 = abc\rangle$, S_4, $P := \langle a, b \mid a^3 = b^8 = e, bab^{-1} = a^{-1}\rangle$ or $Q := \langle a, b \mid a^3 = b^4 = c^2 = e, bab^{-1} = a^{-1}, cbc^{-1} = b^{-1}, ac = ca\rangle$ (One can check it by using the GAP Small Groups Library [1], for example).

By Proposition 6.3, we have $G \not\simeq C_2 \times C_{12}$, $C_2 \times C_2 \times C_6$, Q_{24}, $C_2 \times D_{12}$, $C_2 \times Q_{12}$, $C_2 \times A_4$, $C_3 \times D_8$, $C_3 \times Q_8$, $C_4 \times D_6$, P, Q. Since the orders of maximal cyclic subgroups are $4, 3$ or 2, we have $G \not\simeq C_{24}, D_{24}$. Since $SL(2, 3)$ contains (more than) two non-conjugate cyclic subgroups of order 3, we have $G \not\simeq SL(2, 3)$. The remaining possibility is $G \simeq S_4$.

(iv) $(|G_1|, |G_2|, |G_3|, |G|) = (5, 3, 2, 60)$.

A group of order 60 is isomorphic to C_{60}, $C_2 \times C_{30}$, D_{60}, Q_{60}, $C_3 \times D_{20}$, $C_3 \times Q_{20}$, $C_5 \times D_{12}$, $C_5 \times Q_{12}$, $C_5 \times A_4$, $D_6 \times D_{10}$, A_5, $R := \langle a, b \mid a^{15} = b^4 = e, bab^{-1} = a^2\rangle$ or $S := \langle a, b \mid a^{15} = b^4 = e, bab^{-1} = a^7\rangle$ (One can check it by using the GAP Small Groups Library [1], for example).

By Proposition 6.3, we have $G \not\simeq C_2 \times C_{30}$, $C_3 \times D_{20}$, $C_3 \times Q_{20}$, $C_5 \times D_{12}$, $C_5 \times Q_{12}$, $C_5 \times A_4$, $D_6 \times D_{10}$, R, S. Since the orders of maximal cyclic subgroups of G are $5, 3$ or 2, we have $G \not\simeq C_{60}$, D_{60}, Q_{60}. The remaining possibility is $G \simeq A_5$.

Therefore, the rotational symmetry group of a simple acyclic oriented matroid of rank 4 is one of the following types: C_n $(n \geq 1)$, D_{2n} $(n \geq 1)$, A_4, S_4, A_5.

Next, we prove that all of the above groups can appear as rotational symmetry groups of simple acyclic oriented matroids of rank 4.

(i) C_n.

For $n = 2, 3, \ldots$, let us define 3-dimensional point configurations P_n as follows.

$$\begin{aligned}
P_2 &:= \{(-1,1,0),(0,1,0),(1,1,0),(-1,-1,0),(0,-1,0),(1,-1,0),(0,0,1)\},\\
P_3 &:= \{(\cos(\frac{2k\pi}{3}),\sin(\frac{2k\pi}{3}),-2) \mid k=1,2,3\} \cup \{(\cos(\frac{2k\pi}{3}),\sin(\frac{2k\pi}{3}),2) \mid k=1,2,3\},\\
P_n &:= \{(\cos(\frac{2k\pi}{n}),\sin(\frac{2k\pi}{n}),0) \mid k=1,\ldots,n\} \cup \{(0,0,1)\}\ (n=4,5,\ldots)
\end{aligned}$$

and $\mathcal{M}_{P_n}$ be the oriented matroid associated to P_n. Then $R(\mathcal{M}_{P_n}) \simeq C_n$.

(ii) D_{2n}.
Let $Q_n := \{(\cos(\frac{2k\pi}{n}),\sin(\frac{2k\pi}{n}),0) \mid k=1,\ldots,n\} \cup \{(0,0,-1),(0,0,1)\}$ and $\mathcal{M}_{Q_n}$ be the associated oriented matroid of Q_n. Note that $R(\mathcal{M}_{Q_n}) \supseteq R(Q_n) \simeq D_{2n}$, where $R(Q_n)$ is the geometric rotational symmetry group of Q_n. By the above discussion, $R(\mathcal{M}_{Q_n})$ must be isomorphic to either of C_k $(k \geq 1)$, D_{2k} $(k \geq 1)$, A_4, S_4 or A_5. Simple consideration leads $R(\mathcal{M}_{Q_n}) \simeq D_{2n}$.

(iii) A_4, S_4, A_5.
The oriented matroid associated to the 3-simplex (resp. the 3-cube, Icosahedron) has the rotational symmetry group isomorphic to A_4 (resp. S_4, A_5).



Figure 7: The 3-simplex [left], the 3-cube [center] and Icosahedron [right] contracted by each fixed point

7 Concluding remarks

In this paper, we investigated fixed point properties for oriented matroids. It was shown that some of fixed point properties of geometric rotational symmetry can naturally be extended to the setting of oriented matroids. Using the fixed point theorems, we classified rotational symmetry groups of simple acyclic oriented matroids of rank 4. It turned out that rotational symmetry of oriented matroids of rank $r(\leq 4)$ is similar to that of point configurations. In fact, we showed that the axioms of oriented matroids are powerful enough to develop the theory similar to the classical theory of the classification of finite subgroups of $SO(3)$. Our results imply the possibility that combinatorial behaviors of point configurations are fundamental factors for determining possible symmetry groups of them.

As future work, it may be interesting to investigate the following topics.

- Classification of (full) symmetry groups of simple acyclic oriented matroids of rank 4.
- Classification of (rotational) symmetry groups of simple acyclic oriented matroids of higher ranks.
- Can the existence theorem of fixed points be proved for rotational symmetries of *non-convex* oriented matroids?

- Is there any simple acyclic oriented matroid of rank r whose symmetry group is not isomorphic to any finite subgroups of the orthogonal group $O(r-1)$?

References

[1] H. U. Besche, B. Eick and E. O'Brien, *The Small Groups Library*, `http://www.icm.tu-bs.de/ag_algebra/software/small/`

[2] A. Björner, M. Las Vergnas B. Sturmfels, N. White and G. Ziegler, *Oriented Matroids*, Cambridge University Press, 1999.

[3] R. Bland and M. Las Vergnas, Orientability of matroids, *Journal of Combinatorial Theory Series B* 24:94-123, 1978.

[4] J. Bokowski, G. Ewald and P. Kleinschmidt, On combinatorial and affine automorphisms of polytopes, *Israel J. Math.* 47:123-130, 1984.

[5] R. Cordovil and P. Duchet, On the sign-invariance graphs of uniform oriented matroids, *Discrete Math.*, 79:251-257, 1990.

[6] R. Cordovil and P. Duchet, Cyclic Polytopes and Oriented Matroids, *Europ. J. Combinatorics*, 21:49-64, 2000.

[7] B. Grünbaum, *Convex Polytopes 2nd edition*, Springer, 2003.

[8] J. E. Humphreys, *Reflection groups and Coxeter groups*, volume 29 of Cambridge Studies in Advanced Mathematics, Cambridge University Press, Cambridge, 1990.

[9] M. Las Vergnas, Extensions ponctuelles d'une géométrie combinatoire orienté dans: Problèmes combinatoires et Théorie des graphes, Actes du colloque International CNRS, No. 260, Orsay 1976 (Paris, 1978), pp. 263-268.

[10] P. Mani, Automorphismen von polyedrischen Graphen, *Mathematische Annalen* 192:279-303, 1971.

[11] A. Paffenholz, New Polytopes from Products, *Journal of Combinatorial Theory, Series A* 13:1396-1418, 2006.

[12] J. Richter-Gebert, Two interesting oriented matroids, *Documenta Mathematica* 1:137-148, 1996.

[13] P. W. Shor, Stretchability of pseudolines is NP-hard, *Applied Geometry and Discrete Mathemaics The Victor Klee Festschrift (P. Gritzmann, B. Sturmfels, eds.), DIMACS Series in Discrete Mathematics and Theoretical Computer Science, Amer. Math. Soc., Providence, RI*, 4, 531-554, 1991.

[14] S. Sternberg, *Group Theory and Physics*, Cambridge University Press, 1994.

Appendix: properties related to rank 3 alternating matroids

Here, we give proofs of the propositions related to rank 3 alternating matroids that were not proved above.

Proposition 7.1 Let $\mathcal{M}$ be a simple oriented matroid of rank 3 on a ground set E. Suppose that $\mathcal{M}|_X$ is a relabeling of the alternating matroid $A_{3,|X|}$ for $X \subset E$ and that $x_0, \dots, x_{p-1} \in X$ defines an alternating matroid in this order, i.e., $\chi(x_i, x_j, x_k) = +$ for all $0 \leq i < j < k \leq p-1$, where χ is a chirotope of $\mathcal{M}$ and $p := |X|$. For $i = 0, \dots, p$, let V_i be the cocircuit such that

$$V_i(x_i) = V_i(x_{i+1}) = 0, V_i(x) = + \text{ for all } x \in X \setminus \{x_i, x_{i+1}\},$$

where $x_p := x_0$. Let us consider $e \in E \setminus X$. Then, the sign sequence $V_0(e), V_1(e), \ldots, V_p(e)$ must be one of the following forms:

$$+\cdots+-\cdots-+\cdots+,\ +\cdots+0-\cdots-+\cdots+,\ +\cdots+-\cdots-0+\cdots+,\ +\cdots+0-\cdots-0+\cdots+,$$

$$-\cdots-+\cdots+-\cdots-,\ -\cdots-0+\cdots+-\cdots-,\ -\cdots-+\cdots+0-\cdots-,\ -\cdots-0+\cdots+0-\cdots-,$$

where $+\cdots+$ and $-\cdots-$ may be empty.

Proof: In this proof, we only consider the case where $V_i(e) \neq 0$ for all $i = 0, \ldots, p$. The following discussion can be applied to the other case by a slight modification.

We prove by contradiction. Let us assume that there exist $0 \leq l_1 < l_2 < l_3 \leq p-1$ such that

$$V_{l_1}(e) = +, V_{l_1+1}(e) = -, V_{l_2}(e) = -, V_{l_2+1}(e) = +, V_{l_3}(e) = +, V_{l_3+1}(e) = -.$$

Let W_1, W_2, W_3 be the cocircuits obtained by applying vector elimination to V_{l_1}, V_{l_1+1}, e and V_{l_2}, V_{l_2+1}, e and V_{l_3}, V_{l_3+1}, e, respectively. Then, the cocircuit W_i satisfies

$$W_i(e) = W_i(x_{l_i+1}) = 0,\ W_i(x) = + \text{ for all } x \in X \setminus \{x_{l_i+1}\},$$

for $i = 1, 2, 3$. Apply vector elimination to $W_1, -W_3$ and x_{l_2+1}. Then, we obtain a cocircuit W_4 satisfying

$$W_4(e) = W_4(x_{l_2+1}) = 0, W_4(x_{l_1+1}) = -, W_4(x_{l_3+1}) = +,$$

which is a contradiction (compare with W_2). A contradiction also occurs if there exist $0 \leq l'_1 < l'_2 < l'_3 \leq p-1$ such that

$$V_{l'_1}(e) = -, V_{l'_1+1}(e) = +, V_{l'_2}(e) = +, V_{l'_2+1}(e) = -, V_{l'_3}(e) = -, V_{l'_3+1}(e) = +.$$

□

Proposition 7.2 Let $x_0, \ldots, x_{p-1}$ be the elements of $A_{3,p}$ that define an alternating matroid in this order. For a covector V of $A_{3,p}$ with $V(x_0) = 0$, the sign sequence $V(x_1), V(x_2), \ldots, V(x_{p-1})$ must be one of the following forms:

$$-\cdots-+\cdots+,\ 0-\cdots-+\cdots+,\ -\cdots-0+\cdots+,\ -\cdots-+\cdots+0,$$

$$+\cdots+-\cdots-,\ 0+\cdots+-\cdots-,\ +\cdots+0-\cdots-,\ +\cdots+-\cdots-0,$$

where $+\cdots+$ and $-\cdots-$ may be empty.

Proof: Let us assume that there exist $0 \leq l_1 < l_2 < l_3 \leq p-1$ such that $V(x_{l_1}) = +, V(x_{l_2}) = -, V(x_{l_3}) = +$. Let W be the cocircuit of $A_{3,p}$ such that

$$W(x_0) = W(x_{l_2}) = 0, W(x_i) = + \text{ for } i = 1, \ldots, l_2 - 1, W(x_j) = - \text{ for } j = l_2 + 1, \ldots, p-1$$

and consider vector elimination of V, W and x_{l_3}. Then we obtain a cocircuit Z such that

$$Z(x_0) = Z(x_{l_3}) = 0, Z(x_{l_1}) = +, Z(x_{l_2}) = -,$$

a contradiction. Similarly, we obtain a contradiction if there exist $0 \leq l'_1 < l'_2 < l'_3 \leq p-1$ such that $V(x_{l'_1}) = -, V(x_{l'_2}) = +, V(x_{l'_3}) = -$. □